

# MCS Classicare 2025 Formulary 2 Prior Authorization Criteria

MCS Classicare Del Caribe (HMO), MCS Classicare En Tu Hogar (HMO), MCS Classicare Efectivo (HMO), MCS Classicare Essential (HMO-POS), MCS Classicare Exacto (HMO), MCS Classicare Excede (HMO), MCS Classicare Firme (HMO), MCS Classicare Hero (HMO), MCS Classicare Intelicare (HMO), MCS Classicare MA-PD Group (HMO-POS), MCS Classicare Primero (HMO C-SNP) MCS Classicare Platino Ideal (HMO D-SNP), MCS Classicare Platino MasCa\$h (HMO D-SNP), MCS Classicare Platino Maximo (HMO D-SNP), MCS Classicare Platino Total (HMO D-SNP).

MCS Classicare requires you (or your physician) to get prior authorization for certain drugs. This means that you will need to get approval from MCS Classicare before you fill your prescriptions. If you don't get approval, MCS Classicare may not cover the drug.

MCS Classicare is an HMO plan subscribed by MCS Advantage, Inc.

ATTENTION: If you speak English, language assistance services, free of charge, are available to you. Call 1.866.627.8183 (TTY: 1.866.627.8182).

ATENCIÓN: Si habla español, tiene a su disposición servicios gratuitos de asistencia lingüística. Llame al 1.866.627.8183 (TTY: 1.866.627.8182).

注意:如果您使用繁體中文,您可以免費獲得語言援助服務。請致電 1.866.627.8183 (TTY: 1.866.627.8182).



Medicare Platino Logo: Authorized by the Office of the Election Comptroller OCE-SA 2024-01715

Last Updated: 09/26/2024

### **Prior Authorization Requirements**

## **ABIRATERONE**

### **Products Affected**

• abiraterone acetate oral tablet 250 mg, 500 mg

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer (CRPC), or B.) High risk, castration-sensitive metastatic prostate cancer (CSPC). For treatment of CRPC and CSPC, abiraterone will be used in combination with prednisone AND one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog (e.g. leuprolide, triptorelin), OR 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **ACITRETIN**

### **Products Affected**

• acitretin

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Severely impaired liver or kidney function, B.) Chronic abnormally elevated blood lipid values, or C.) Concomitant use of methotrexate or tetracyclines
Required Medical Information	Diagnosis of severe, recalcitrant psoriasis (including plaque, guttate, erythrodermic palmar- plantar and pustular) AND patient must have tried and failed, contraindication or intolerance to one formulary first line agent (e.g., Topical Corticosteroids (betamethasone, fluocinonide, desoximetasone), Topical Calcipotriene/Calcitriol, Topical Calcipotriene, OR Topical Tazarotene)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a dermatologist
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ACTEMRA-2**

### **Products Affected**

ACTEMRA

### • ACTEMRA ACTPEN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Giant cell arteritis, B.) Polyarticular juvenile idiopathic arthritis (PJIA) and patient has trial and failure or intolerance or contraindication to two preferred products, (i.e. Adalimumab-ADBM/Humria, Enbrel), C.) Rheumatoid arthritis (RA) and patient has trial and failure or intolerance or contraindication to two preferred products, (i.e. Adalimumab-ADBM/Humira, Enbrel, Rinvoq), D.) Systemic juvenile idiopathic arthritis (SJIA), or E.) Systemic sclerosis-associated interstitial lung disease
Age Restrictions	2 years of age and older
Prescriber Restrictions	For RA, pJIA: Rheumatologist
Coverage Duration	12 months
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **ACTIMMUNE**

### **Products Affected**

• ACTIMMUNE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections, or B.) Severe, malignant osteopetrosis (SMO)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ADALIMUMAB-ADBM**

#### **Products Affected**

- adalimumab-adbm (2 pen)
- adalimumab-adbm (2 syringe) subcutaneous prefilled syringe kit 10 mg/0.2ml, 20 mg/0.4ml, 40 mg/0.8ml
- adalimumab-adbm(cd/uc/hs strt)
- adalimumab-adbm(ps/uv starter)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Noninfectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa. For moderately to severely active rheumatoid arthritis: 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX), or corticosteroids. For active ankylosis spondylitis: patient has experienced an inadequate treatment response, intolerance, or contraindication to a non-steroidal anti-inflammatory drug(NSAID). For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	For Juvenile Idiopathic Arthritis: 2 years of age or older. For Pediatric Crohn's disease: 6 years of age or older. For Hidradenitis Suppurativa: 12 years of age or older. For Uveitis: 2 years of age or older. For Ulcerative

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
	Colitis: 5 years of age or older. All other indications 18 years of age or older.
Prescriber Restrictions	1) For PsO and HS: Dermatologist, 2) For CD and UC: Gastroenterologist, 3) For Uveitis: Ophthalmologist, 4) For RA, PJIA, and AS: Rheumatologist, 5) For PsA: Dermatologist or Rheumatologist.
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ADEMPAS**

### **Products Affected**

• ADEMPAS

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **AFINITOR**

#### **Products Affected**

• everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Hypersensitivity to everolimus, or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus)
Required Medical Information	Diagnosis of one of the following A.) Renal angiomyolipoma and tuberous sclerosis complex (TSC) not requiring immediate surgery, B.) Advanced hormone receptor-positive, HER2 negative breast cancer in postmenopausal women and taken in combination with exemestane, after failure with letrozole or anastrozole, C.) Progressive, well-differentiated, nonfunctional neuroendocrine tumors of gastrointestinal or lung origin and disease is unresectable, locally advanced, or metastatic, D.) Pancreatic progressive neuroendocrine tumors and disease is unresectable, locally advanced, or metastatic, E.) Advanced renal cell carcinoma (RCC) after failure with sunitinib or sorafenib, F.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **AKEEGA**

### **Products Affected**

AKEEGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of deleterious or suspected deleterious BRCA-mutated (BRCAm) metastatic castration-resistant prostate cancer (mCRPC) as detected on an FDA-approved test AND used in combination with prednisone
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ALECENSA**

### **Products Affected**

ALECENSA

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **ALOSETRON**

### **Products Affected**

• alosetron hcl

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Constipation, B.) History of Chronic or severe constipation or sequelae from constipation, C.) History of ischemic colitis, intestinal obstruction, stricture, toxic megacolon, GI perforation, adhesions, diverticulitis, Crohn's disease, ulcerative colitis, D.) History of severe hepatic impairment, E.) History of impaired intestinal circulation, thrombophlebitis, or hypercoagulable state, or F.) Coadministration with fluvoxamine
Required Medical Information	Diagnosis of irritable bowel syndrome, severe diarrhea-predominant
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ALUNBRIG**

### **Products Affected**

• ALUNBRIG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase-positive (ALK) metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **AMBRISENTAN**

### **Products Affected**

• ambrisentan

PA Criteria	Criteria Details
Exclusion Criteria	Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension
Required Medical Information	Diagnosis of pulmonary arterial hypertension classified as WHO Group I, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	12 months
Other Criteria	Continuation of therapy: patient is tolerating and responding to medication and has improved exercise capacity or a delay in clinical worsening.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **APTIOM**

### **Products Affected**

• APTIOM

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of partial-onset seizures
Age Restrictions	4 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ARCALYST**

### **Products Affected**

ARCALYST

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **ARIKAYCE**

### **Products Affected**

• ARIKAYCE

PA Criteria	Criteria Details
Exclusion Criteria	Known sensitivity to any aminoglycoside
Required Medical Information	Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients (greater than 6 months of a multidrug background regimen)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist or pulmonologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ATYPICALS**

#### **Products Affected**

- CAPLYTA
- FANAPT
- FANAPT TITRATION PACK
- LYBALVI
- REXULTI

- SECUADO
- VERSACLOZ
- VRAYLAR ORAL CAPSULE 1.5 MG, 3 MG, 4.5 MG, 6 MG

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **AUGTYRO**

### **Products Affected**

• AUGTYRO

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **AUSTEDO**

#### **Products Affected**

- AUSTEDO
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HOUR 12 MG, 18 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR PATIENT TITRATION ORAL TABLET EXTENDED RELEASE THERAPY PACK 12 & 18 & 24 & 30 MG, 6 & 12 & 24 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression in a patient with Huntington's Disease, B.) Hepatic impairment, C.) Concomitant use of MAOIs, reserpine, tetrabenazine, or valbenazine
Required Medical Information	Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist or psychiatrist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **AUVELITY**

### **Products Affected**

AUVELITY

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **AYVAKIT**

### **Products Affected**

AYVAKIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations, B.) Advanced Systemic Mastocytosis (AdvSM), including aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SMAHN), or mast cell leukemia (MCL), and platelet count of at least 50,000/mcL, or C.) Indolent systemic mastocytosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, allergist, or immunologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BALVERSA**

### **Products Affected**

• BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **BANZEL**

### **Products Affected**

• rufinamide oral tablet

PA Criteria	Criteria Details
Exclusion Criteria	Familial Short QT Syndrome
Required Medical Information	Diagnosis of seizures associated with Lennox-Gastaut syndrome
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BENLYSTA**

### **Products Affected**

• BENLYSTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Active, autoantibody-positive, system lupus erythematosus (SLE), or B.) Active lupus nephritis and patient is receiving standard therapy (corticosteroids, antimalarials, NSAIDs, immunosuppressives)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist or rheumatologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BESREMI**

### **Products Affected**

• BESREMI

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Existence of, or history of severe psychiatric disorders (severe depression, suicidal ideation, or suicide attempt), B.) Hypersensitivity to interferons including interferon alfa-2b or excipients, C.) Hepatic impairment (Child-Pugh B or C), D.) History or presence of active serious or untreated autoimmune disease, or E.) Immunosuppressed transplant recipients
Required Medical Information	Diagnosis of polycythemia vera
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BEXAROTENE GEL**

### **Products Affected**

• bexarotene

	T
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL Stage 1A/1B) and patient had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) indicated for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or dermatologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BEXAROTENE ORAL**

### **Products Affected**

• bexarotene

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, hematologist, or dermatologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BOSENTAN**

### **Products Affected**

• bosentan

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant cyclosporine A or glyburide therapy
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I) and patient has New York Heart Association (NYHA) Functional Class II-IV, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **BOSULIF**

#### **Products Affected**

• BOSULIF ORAL CAPSULE 100 MG, 50 • BOSULIF ORAL TABLET 100 MG, 400 MG, 500 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic, accelerated, or blast phase Philadelphia chromosome-positive (Ph+) chronic myelogenous leukemia (CML) with resistance or inadequate response to prior therapy, or B.) Newly diagnosed chronic phase Philadelphia chromosome-positive (Ph+) CML
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BRAFTOVI**

### **Products Affected**

• BRAFTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) unresectable or metastatic melanoma with documented BRAF V600E or V600K mutation as detected by a FDA-approved test and used in combination with binimetinib, B.) metastatic colorectal cancer with documented BRAF V600E mutation as detected by a FDA-approved test, patient has received prior therapy, and braftovi used in combination with cetuximab, or C.) Metastatic non-small cell lung cancer with a BRAF V600E mutation as detected by an FDA-approved test and used in combination with binimetinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BRIVIACT**

### **Products Affected**

• BRIVIACT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of partial-onset seizures, member must have history of inadequate response, contraindication, or intolerance to levetiracetam prior to approval.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **BRONCHITOL**

### **Products Affected**

• BRONCHITOL

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **BRUKINSA**

### **Products Affected**

• BRUKINSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) mantle cell lymphoma (MCL) and patient has received at least one prior therapy, B.) Treatment of adult patients with Waldenstrom macroglobulinemia, C.) Treatment of adult patients with relapsed or refractory marginal zone lymphoma who have received at least one anti-CD20-based regimen (e.g., rituximab, obinutuzumab, ofatumumab), D.) Chronic lymphocytic leukemia, E.) Small lymphocytic lymphoma, or F.) Relapsed or refractory follicular lymphoma, in combination with obinutuzumab, after 2 or more lines of systemic therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **CABOMETYX**

### **Products Affected**

CABOMETYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Advanced hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib, C.) Advanced renal cell carcinoma and used as first line treatment in combination with nivolumab or D.) treatment of adults and pediatric patients 12 years and older with locally advanced or metastatic differentiated thyroid cancer that has progressed following VEGFR-targeted therapy (e.g., Axitinib, Lenvatinib, Pazopanib, Sorafenib, Sunitinib, vandetanib) and who are radioactive iodine-refractory or ineligible
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **CALQUENCE**

### **Products Affected**

• CALQUENCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least 1 prior therapy, B.) Chronic lymphocytic leukemia (CLL), or C.) Small lymphocytic lymphoma (SLL)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **CAMZYOS**

#### **Products Affected**

• CAMZYOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) in adult patients
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **CAPRELSA**

#### **Products Affected**

• CAPRELSA ORAL TABLET 100 MG, 300 MG

PA Criteria	Criteria Details
Exclusion Criteria	Congenital long QT syndrome
Required Medical Information	Diagnosis of metastatic or unresectable locally advanced medullary thyroid cancer (MTC) AND disease is symptomatic or progressive
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **CARGLUMIC ACID**

#### **Products Affected**

• carglumic acid

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) N-acetyl glutamate synthase (NAGS) deficiency (confirmed by appropriate genetic testing) with acute or chronic hyperammonemia, or B.) Propionic or methylmalonic acidemia with acute hyperammonemia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **CAYSTON**

#### **Products Affected**

• CAYSTON

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **CIMZIA-2**

#### **Products Affected**

• CIMZIA

• CIMZIA (2 SYRINGE)

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **CLOBAZAM**

#### **Products Affected**

- clobazam oral suspension
- clobazam oral tablet

#### SYMPAZAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of seizures associated with Lennox-Gastaut syndrome
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **CNS STIMULANTS**

#### **Products Affected**

• armodafinil

• modafinil

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work disorder (SWD)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **COMETRIQ**

#### **Products Affected**

- COMETRIQ (100 MG DAILY DOSE)
  - COMETRIQ (140 MG DAILY DOSE)
- COMETRIQ (60 MG DAILY DOSE)

COMETRIQ (140 MO DAILT DOSE)	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of progressive, metastatic medullary thyroid cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **COPIKTRA**

#### **Products Affected**

COPIKTRA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory chronic lymphocytic leukemia (CLL), or B.) Relapsed or refractory small lymphocytic lymphoma (SLL). For CLL or SLL, the patient must have history of at least 2 prior therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **COSENTYX**

#### **Products Affected**

- COSENTYX (300 MG DOSE)
- COSENTYX UNOREADY
- COSENTYX SENSOREADY (300 MG)

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **COTELLIC**

#### **Products Affected**

• COTELLIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.)unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf), or B.) Histiocytic neoplasms
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **CYSTAGON**

#### **Products Affected**

• CYSTAGON

PA Criteria	Criteria Details
Exclusion Criteria	Known serious hypersensitivity to penicillamine or cysteamine
Required Medical Information	Diagnosis of nephropathic cystinosis confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a nephrologist
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DALFAMPRIDINE**

#### **Products Affected**

• dalfampridine er

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute)
Required Medical Information	Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DAURISMO**

#### **Products Affected**

DAURISMO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with cytarabine in patients 75 years of age or older OR in patients that have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an Oncologist or Hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DAYBUE**

#### **Products Affected**

• DAYBUE

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **DEFERASIROX**

#### **Products Affected**

• deferasirox

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10(9)/L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS)
Required Medical Information	Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DEFERIPRONE**

#### **Products Affected**

• deferiprone

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of transfusional iron overload due to thalassemia syndromes, sickle cell disease, or other anemias 2.) failure to current chelation therapy (e.g., deferoxamine)
Age Restrictions	8 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DIACOMIT**

#### **Products Affected**

• DIACOMIT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe myoclonic epilepsy in infancy (Dravet syndrome) in patients taking clobazam
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DICLOFENAC PATCH**

#### **Products Affected**

• diclofenac epolamine

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute mild to moderate pain due to minor strains, sprains and contusions
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DIMETHYL FUMARATE**

#### **Products Affected**

• dimethyl fumarate

• dimethyl fumarate starter pack

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DRONABINOL**

#### **Products Affected**

• dronabinol

PA Criteria	Criteria Details
Exclusion Criteria	Sesame oil hypersensitivity
Required Medical Information	Diagnosis of one of the following A.) Anorexia associated to AIDS, or B.) Chemotherapy-induced nausea and vomiting
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **DROXIDOPA**

#### **Products Affected**

• droxidopa

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Initial: 1 month, Renewal: 3 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **DUPIXENT**

#### **Products Affected**

DUPIXENT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	For atopic dermatitis (AD), initial therapy: 1) Patient has moderate-to-severe disease, AND 2) Patient has had an inadequate treatment response to either a topical corticosteroid or a topical calcineurin inhibitor, OR topical corticosteroids and topical calcineurin inhibitors are not advisable for the patient. For oral corticosteroid dependent asthma, initial therapy: Patient has inadequate asthma control despite current treatment with both of the following medications: 1) High-dose inhaled corticosteroid AND 2) Additional controller (i.e., long acting beta2-agonist, long-acting, muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For moderate-to-severe asthma, initial therapy: Patient has a baseline blood eosinophil count of at least 150 cells per microliter and their asthma remains inadequately controlled despite current treatment with both of the following medications: 1) Medium-to-high-dose inhaled corticosteroid, AND 2) Additional controller (i.e., long acting beta2-agonist, long-acting muscarinic antagonist, leukotriene modifier, or sustained-release theophylline) unless patient has an intolerance or contraindication to such therapies. For asthma, continuation of therapy: Asthma control has improved on treatment with the requested drug, as demonstrated by a reduction in the frequency and/or severity of symptoms and exacerbations or a reduction in the daily maintenance oral corticosteroid dose. For chronic rhinosinusitis with nasal polyposis (CRSwNP): 1) The requested drug is used as add-on maintenance treatment, AND 2) Patient has experienced an inadequate treatment response to fluticasone. For Eosinophilic esophagitis failure, intolerance or contraindication to PPIs
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an allergist, dermatologist, gastroenterologist, immunologist, otolaryngologist, pulmonologist, or rheumatologist

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **EMGALITY-1-2**

#### **Products Affected**

EMGALITY

• EMGALITY (300 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic or episodic migraine disorder and patient has documented trial, inadequate response, or contraindication to at least 2 generic formulary drugs used for migraine prevention (i.e., propranolol, topiramate, divalproex, timolol), or B.) Episodic cluster headache
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **EMSAM**

#### **Products Affected**

• EMSAM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use with any of the following: SSRIs, SNRIs, clomipramine, imipramine, meperidine, tramadol, methadone, pentazocine, propoxyphene, dextromethorphan, carbamazepine, or B.) Pheochromocytoma
Required Medical Information	Diagnosis of major depressive disorder and patient had trial of at least 2 generic oral antidepressants from differing classes (at least one should be from the following list: selective serotonin reuptake inhibitors, serotonin and norepinephrine reuptake inhibitors, mirtazapine, or bupropion unless contraindicated), unless unable to take any oral medication AND Patient had an adequate washout period (for patients previously on agents requiring a washout period)
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ENBREL**

#### **Products Affected**

- ENBREL
- ENBREL MINI

#### • ENBREL SURECLICK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy. For moderately to severely active rheumatoid arthritis: 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX), or corticosteroids. For active ankylosing spondylitis: patient has experienced an inadequate treatment response, intolerance, or contraindication to a nonsteroidal anti-inflammatory drug(NSAID). For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	2 years of age for JIA and PsA, and 4 years of age for PsO
Prescriber Restrictions	1) For RA, PJIA, AS: Rheumatologist, 2) For PsA: Dermatologist or Rheumatologist, 3) For PsO: Dermatologist.
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
	with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ENSPRYNG**

#### **Products Affected**

ENSPRYNG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active Hepatitis B infection, or B.) Active or untreated latent tuberculosis
Required Medical Information	Diagnosis of neuromyelitis optica spectrum disorder (NMOSD) in patients who are anti-aquaporin-4 (AQP4) antibody positive
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist, immunologist, or ophthalmologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **EPIDIOLEX**

#### **Products Affected**

• EPIDIOLEX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Lennox-Gastaut syndrome, B.) Severe myoclonic epilepsy in infancy (Dravet syndrome), or C.) Seizures associated with tuberous sclerosis complex (TSC)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **EPOETIN THERAPY**

### **Products Affected**

• RETACRIT

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **ERIVEDGE**

#### **Products Affected**

• ERIVEDGE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic basal cell carcinoma, or B.) Locally advanced basal cell carcinoma that has recurred following surgery or the patient is not a candidate for surgery or radiation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ERLEADA**

#### **Products Affected**

ERLEADA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Nonmetastatic, castration-resistant prostate cancer (nmCRPC), or B.) Metastatic, castration-sensitive prostate cancer (mCSPC). For treatment of nmCRPC and mCSPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog, OR 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **ERLOTINIB**

#### **Products Affected**

• erlotinib hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib will be used in combination with gemcitabine, B.) Locally advanced or metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND one of the following 1.) Erlotinib will be used as first-line treatment, 2.) Failure with at least one prior chemotherapy regimen, OR 3.) No evidence of disease progression after four cycles of first-line platinum-based chemotherapy and erlotinib will be used as maintenance treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **EVEROLIMUS SUSPENSION**

#### **Products Affected**

• everolimus oral tablet soluble 2 mg, 3 mg, 5 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Hypersensitivity to everolimus, or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus)
Required Medical Information	Diagnosis of one of the following A.) Tuberous sclerosis complex (TSC)-associated partial-onset seizures, or B.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **EVRYSDI**

#### **Products Affected**

EVRYSDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of spinal muscular atrophy (SMA)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FASENRA**

### **Products Affected**

FASENRA

### • FASENRA PEN

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **FEBUXOSTAT**

### **Products Affected**

• febuxostat

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of azathioprine or mercaptopurine
Required Medical Information	Diagnosis of Gout and all of the following 1.) documented inadequate treatment response, adverse event, or contraindication to maximally titrated dose of Allopurinol, and 2.) patients with established cardiovascular disease, prescriber attests that benefit of treatment outweighs the risk of treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## FENTANYL ORAL

### **Products Affected**

• fentanyl citrate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients, C.) Known or suspected gastrointestinal obstruction, including paralytic ileus, D.) Acute or severe bronchial asthma and used in an unmonitored setting (absence of resuscitative equipment)
Required Medical Information	Must meet all of the following 1.) Diagnosis of cancer-related breakthrough pain, 2.) Patient is currently receiving/tolerant to around-the-clock opioid therapy for persistent cancer pain, and 3.) Patient and prescriber are enrolled in the TIRF REMS Access Program
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FILSPARI**

### **Products Affected**

• FILSPARI

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with angiotensin receptor blockers (ARBs), endothelin receptor antagonists (ERAs), or aliskiren
Required Medical Information	Diagnosis of treatment of primary immunoglobulin A (IgA) nephropathy at risk of rapid disease progression, generally a urine protein to creatinine ratio (UPCR) of 1.5 g/g or more, to reduce proteinuria
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FIRMAGON**

### **Products Affected**

• FIRMAGON

• FIRMAGON (240 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FOTIVDA**

### **Products Affected**

• FOTIVDA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory advanced renal cell cancer (RCC) following 2 or more prior systemic therapies
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **FRUZAQLA**

### **Products Affected**

• FRUZAQLA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic colorectal cancer (mCRC) and all of the following: A.) patient has been previously treated with fluoropyrimidine, oxaliplatin, irinotecan-based chemotherapy, B.) an anti-VEGF therapy, and C.) if RAS wild-type and medically appropriate, patient has also been previously treated with anti-EGFR therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FULPHILA**

### **Products Affected**

• FULPHILA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of a non-myeloid malignancy and drug is being used as prophylaxis for chemotherapy-induced neutropenia
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	3 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **FYCOMPA**

### **Products Affected**

FYCOMPA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Partial-onset seizures with or without secondary generalization, or B.) Primary generalized tonic-clonic seizure disorder as adjunctive therapy
Age Restrictions	4 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **GAVRETO**

### **Products Affected**

GAVRETO

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **GEFITINIB**

### **Products Affected**

• gefitinib

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) and must meet all of the following 1.) Tumor has epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility and 2.) Used as first-line treatment
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **GILOTRIF**

### **Products Affected**

• GILOTRIF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC with progression after platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **GLATIRAMER**

### **Products Affected**

• glatiramer acetate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **GLEOSTINE**

### **Products Affected**

• GLEOSTINE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet one of the following: A.) Hodgkin's disease in patient who has relapsed during or failed to respond to primary therapy and is being used in combination with other agents OR B.) Intracranial tumor
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **GLP1**

#### **Products Affected**

- BYDUREON BCISE
- MOUNJARO
- OZEMPIC (0.25 OR 0.5 MG/DOSE)
- OZEMPIC (1 MG/DOSE) SUBCUTANEOUS SOLUTION PEN-INJECTOR 2 MG/1.5ML, 4 MG/3ML
- OZEMPIC (2 MG/DOSE)
- RYBELSUS
- TRULICITY

PA Criteria	Criteria Details
Exclusion Criteria	Used for weightloss
Required Medical Information	The drug is prescribed for an FDA-approved indication
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **GROWTH HORMONE**

### **Products Affected**

OMNITROPE

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Use for growth promotion in pediatric patients with closed epiphyses, B.) Acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) Active malignancy, D.) Active proliferative or severe nonproliferative diabetic retinopathy, E.) Prader-Willi Syndrome in patients who are severely obese, have a history of upper airway obstruction or sleep apnea, or have severe respiratory impairment
Required Medical Information	Supporting statement of diagnosis from the physician
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist or nephrologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HEPATITIS B**

#### **Products Affected**

- adefovir dipivoxil
- BARACLUDE

- entecavir
- VEMLIDY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic hepatitis B and all of the following 1.) Patient has or had evidence of viral replication prior to initiation, 2.) Patient has evidence of persistent elevations in serum aminotransferase (ALT or AST) or histologically active disease, and 3.) Patient is receiving anti-retroviral therapy if the patient has HIV co-infection
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HEPATITIS C**

#### **Products Affected**

- ledipasvir-sofosbuvir
- MAVYRET

• sofosbuvir-velpatasvir

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must submit documentation of chronic hepatitis C genotype (confirmed by HCV RNA level within the last 6 months). Must submit laboratory results within 12 weeks prior to initiating therapy including: 1) CBC w Platelets, 2) AST/ALT, 3) Total Bilirubin, 4) Serum Albumin, and 5) PT/INR. Genotype and subtype are not required for: (1) initial treatment of patients without cirrhosis if using Sofosbuvir-Velpatasvir or Mavyret OR (2) treatment of patients with compensated cirrhosis if using Mavyret
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist
Coverage Duration	Duration of approval per AASLD Guidelines
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANALGESICS**

### **Products Affected**

• butalbital-acetaminophen

• butalbital-apap-caffeine

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANTI-ARRHYTHMICS**

### **Products Affected**

- DIGOXIN ORAL SOLUTION
- digoxin oral tablet 250 mcg

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary Non HRM alternatives: propranolol, sotalol, dofetilide, amiodarone, propafenone, mexiletine
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANTICHOLINERGIC**

### **Products Affected**

• chlordiazepoxide-clidinium

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANTIDEMENTIA**

### **Products Affected**

• ergoloid mesylates

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary non-HRM alternatives: donepezil, galantamine, memantine, rivastigmine oral
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANTIDEPRESSANTS**

### **Products Affected**

• chlordiazepoxide-amitriptyline

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary non-HRM alternatives: mirtazapine, trazodone, fluoxetine, escitalopram, fluvoxamine, desvenlafaxine, duloxetine, sertraline, venlafaxine, bupropion
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - ANTIHISTAMINES**

#### **Products Affected**

- hydroxyzine hcl
- hydroxyzine pamoate

- promethazine hcl
- promethegan

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary non-HRM alternatives FOR MANAGEMENT OF ALLERGIC CONDITIONS: deslorated ine, cetirizine syp, levocetirizine. FOR MANAGEMENT OF ANXIETY/SEDATION: buspirone, trazodone. FOR MANAGEMENT OF NAUSEA/VOMITING: granisetron, ondansetron
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **HRM - ANTIPARKINSON AGENTS**

### **Products Affected**

• benztropine mesylate

• trihexyphenidyl hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary non-HRM alternatives: amantadine, carbidopa/levodopa, entacapone, Neupro, pramipexole, ropinirole, selegiline
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - BENZODIAZEPINES**

#### **Products Affected**

- alprazolam oral tablet 0.25 mg, 0.5 mg, 1 estazolam mg, 2 mg
  - temazepam

• chlordiazepoxide hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) The prescribing physician attests to the medical necessity for using this high risk medication. For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs)
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Part B Prerequisite	No

## **HRM - CARDIOVASCULAR**

### **Products Affected**

• guanfacine hcl er

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary Non HRM alternatives: atomoxetine, dexmethylphenidate, methylphenidate
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - MEGESTROL**

### **Products Affected**

megestrol acetate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication.
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary Non HRM alternatives: dronabinol
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HRM - PCD - BENZODIAZEPINES**

#### **Products Affected**

- clonazepam oral tablet 0.5 mg, 1 mg, 2 mg •
- clonazepam oral tablet dispersible 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg
- clorazepate dipotassium
- diazepam intensol

- diazepam oral solution
- diazepam oral tablet
- lorazepam
- lorazepam intensol

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of seizure disorder or must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) The prescribing physician attests to the medical necessity for using this high risk medication. For all indications: The prescriber must acknowledge the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.) For the management of anxiety disorders: 1) The requested drug is being used concurrently with a selective serotonin reuptake inhibitor (SSRI) or serotonin-norepinephrine reuptake inhibitor (SNRI) until the SSRI/SNRI becomes effective for the symptoms of anxiety, OR 2) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to AT LEAST TWO agents from the following classes: a) selective serotonin reuptake inhibitors (SSRIs), b) serotonin-norepinephrine reuptake inhibitors (SNRIs).
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **HRM - SEDATIVE HYPNOTICS**

### **Products Affected**

• zaleplon

• zolpidem tartrate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) The patient has experienced an inadequate treatment response, intolerance, or has a contraindication to doxepin (3 mg or 6 mg)3.) The prescriber must acknowledge that the benefit of therapy with this prescribed medication outweighs the potential risks for the patient. (Note: The American Geriatrics Society identifies the use of this medication as potentially inappropriate in older adults, meaning it is best avoided, prescribed at reduced dosage, or used with caution or carefully monitored.)
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older.
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Formulary non-HRM alternatives: Belsomra, doxepin 3mg and 6mg tablets, trazodone
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

# HRM - SKELETAL MUSCLE RELAXANTS

#### **Products Affected**

• cyclobenzaprine hcl

• orphenadrine citrate er

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) The drug is being prescribed for an FDA-approved indication, 2.) Patient has tried and failed at least one non-HRM alternative, 3.) The prescribing physician attests to the medical necessity for using this high risk medication
Age Restrictions	Automatic approval if member is less than 65 years of age. Prior Auth required for age 65 or older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	Formulary Non HRM alternatives: baclofen
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **HUMIRA**

#### **Products Affected**

- HUMIRA (2 PEN) SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.4ML, 40 MG/0.8ML, 80 MG/0.8ML
- HUMIRA (2 SYRINGE)
   SUBCUTANEOUS PREFILLED
   SYRINGE KIT 10 MG/0.1ML, 20
- MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA-CD/UC/HS STARTER
- HUMIRA-PED>/=40KG UC STARTER
- HUMIRA-PSORIASIS/UVEIT STARTER

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Noninfectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa. For moderately to severely active rheumatoid arthritis: 1) patient has experienced an inadequate treatment response, intolerance, or has a contraindication to methotrexate (MTX), or corticosteroids. For active ankylosis spondylitis: patient has experienced an inadequate treatment response, intolerance, or contraindication to a non-steroidal anti-inflammatory drug(NSAID). For moderate to severe plaque psoriasis (new starts only): 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin, is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected)

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Age Restrictions	For Juvenile Idiopathic Arthritis: 2 years of age or older. For Pediatric Crohn's disease: 6 years of age or older. For Hidradenitis Suppurativa: 12 years of age or older. For Uveitis: 2 years of age or older. For Ulcerative Colitis: 5 years of age or older. All other indications 18 years of age or older.
Prescriber Restrictions	1) For PsO and HS: Dermatologist, 2) For CD and UC: Gastroenterologist, 3) For Uveitis: Ophthalmologist, 4) For RA, PJIA, and AS: Rheumatologist, 5) For PsA: Dermatologist or Rheumatologist.
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **HYFTOR**

### **Products Affected**

• HYFTOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Facial angiofibroma associated with tuberous sclerosis
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **IBRANCE**

#### **Products Affected**

• IBRANCE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with fulvestrant and disease has progressed following endocrine therapy, or B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and used in combination with an aromatase inhibitor in a male or female patient as initial endocrine-based therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **ICATIBANT**

#### **Products Affected**

• icatibant acetate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hereditary angioedema AND medication will be used for the treatment of acute attacks
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an allergist, hematologist, or immunologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **ICLUSIG**

#### **Products Affected**

• ICLUSIG ORAL TABLET 10 MG, 15 MG, 30 MG, 45 MG

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **IDHIFA**

#### **Products Affected**

• IDHIFA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase 2 (IDH2) mutation as detected by an FDA approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **IMATINIB**

#### **Products Affected**

• imatinib mesylate oral tablet 100 mg, 400 mg

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **IMBRUVICA**

#### **Products Affected**

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, C.) Waldenstrom's macroglobulinemia (WM), or D.) Chronic graft vs host disease (cGVHD) after failure of at least one first-line corticosteroid therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **IMPAVIDO**

#### **Products Affected**

• IMPAVIDO

PA Criteria	Criteria Details
Exclusion Criteria	Sjogren-Larsson-Syndrome
Required Medical Information	Diagnosis of one of the following: A.) Visceral leishmaniasis caused by Leishmania donovani, B.) Cutaneous leishmaniasis caused by Leishmania braziliensis, Leishmania guyanensis, and Leishmania panamensis, or C.) Mucosal leishmaniasis caused by Leishmania braziliensis
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **INCRELEX**

#### **Products Affected**

INCRELEX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Active or suspected malignancy, B.) Use for growth promotion in patients with closed epiphyses, or C.) Intravenous administration
Required Medical Information	Diagnosis of one of the following A.) Severe primary insulin-like growth factor-1 (IGF-1) deficiency and utilized for pediatric treatment of growth failure, or B.) Growth hormone (GH) gene deletion and patient has developed neutralizing antibodies to GH and utilized for pediatric treatment of growth failure
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### INJECTABLE TESTOSTERONE

#### **Products Affected**

• testosterone cypionate

• testosterone enanthate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Carcinoma of the breast (males only), or B.) Known or suspected carcinoma of the prostate
Required Medical Information	Diagnosis of one of the following A.) Hypogonadotropic hypogonadism, B.) Inoperable metastatic breast cancer in women who are postmenopausal (testosterone enanthate), C.) Primary hypogonadism, or D.) Delayed puberty (testosterone enanthate). Diagnosis of hypogonadism must be confirmed by a low-for-age serum testosterone (total or free) level defined by the normal laboratory reference value
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **INLYTA**

#### **Products Affected**

INLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma and patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens), or B.) Advanced renal cell carcinoma and used as first-line therapy in combination with avelumab or pembrolizumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **INQOVI**

#### **Products Affected**

• INQOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo and secondary MDS with the following French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, and chronic myelomonocytic leukemia [CMML]) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **INREBIC**

#### **Products Affected**

• INREBIC

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF).
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **INVEGA HAFYERA**

#### **Products Affected**

• INVEGA HAFYERA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 1092 MG/3.5ML, 1560 MG/5ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following A.) Diagnosis of schizophrenia, and B.) Adequate treatment has been established with Invega Sustenna for at least 4 months or Invega Trinza for at least one three-month injection cycle
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an psychiatrist
Coverage Duration	Plan Year
Other Criteria	For continuation of therapy member must be re-evaluated by the psychiatrist to assess clinical response. Treatment should be discontinued if failure to therapy or an adverse event is documented or the member does not demonstate adherence to treatment
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **INVEGA SUSTENNA**

#### **Products Affected**

 INVEGA SUSTENNA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE 117 MG/0.75ML, 156 MG/ML, 234 MG/1.5ML, 39 MG/0.25ML, 78 MG/0.5ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet one of the following A.) Diagnosis of schizophrenia and all of the following 1.) trial and failure, contraindication, or intolerance to two oral second-generation antipsychotics, 2.) Trial and failure to respond to haloperidol decanoate and fluphenazine decanoate, 3.)Member has established response and tolerability to oral paliperidone, and 4.) Patient presents history of noncompliance with oral medications and documentation is provided on the efforts to improve medication adherence, or B.) Diagnosis of schizoaffective disorder and both of the following 1.)Member has established response and tolerability to oral paliperidone and 2.) Patient presents history of noncompliance with oral medications and documentation is provided on the efforts to improve medication adherence
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an psychiatrist
Coverage Duration	Plan Year
Other Criteria	For continuation of therapy member must be re-evaluated by the psychiatrist to assess clinical response. Treatment should be discontinued if failure to therapy or an adverse event is documented or the member does not demonstate adherence to treatment
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **INVEGA TRINZA**

#### **Products Affected**

• INVEGA TRINZA INTRAMUSCULAR SUSPENSION PREFILLED SYRINGE

273 MG/0.88ML, 410 MG/1.32ML, 546 MG/1.75ML, 819 MG/2.63ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following A.) Diagnosis of schizophrenia, and B.) Adequate treatment has been established with Invega Sustenna for at least 4 months
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an psychiatrist
Coverage Duration	Plan Year
Other Criteria	For continuation of therapy member must be re-evaluated by the psychiatrist to assess clinical response. Treatment should be discontinued if failure to therapy or an adverse event is documented or the member does not demonstate adherence to treatment
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **ITRACONAZOLE**

#### **Products Affected**

• itraconazole

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **IVABRADINE**

#### **Products Affected**

• ivabradine hcl

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **IVERMECTIN**

#### **Products Affected**

• ivermectin

PA Criteria	Criteria Details
Exclusion Criteria	Prevention or treatment of COVID-19
Required Medical Information	Diagnosis of one of the following: A.) Strongyloidiasis of the intestinal tract or B.) Onchocerciasis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **IWILFIN**

#### **Products Affected**

• IWILFIN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of high-risk neuroblastoma to be used to reduce the risk of relapse in adult and pediatric patients who have demonstrated at least a partial response to prior multiagent, multimodality therapy including anti-GD2 immunotherapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **JAKAFI**

#### **Products Affected**

JAKAFI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis, B.) Polycythemia vera AND patient has had an inadequate response to or is intolerant of hydroxyurea, C.) Acute graft versus host disease AND disease is refractory to steroid therapy, or D.) Chronic graft-versus-host disease (cGVHD) after failure of corticosteroid therapy (alone or in combination with a calcineurin inhibitor) and up to one additional line of systemic therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **JAYPIRCA**

#### **Products Affected**

• JAYPIRCA ORAL TABLET 100 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) relapsed or refractory mantle cell lymphoma (MCL) and is being used after at least two lines of systemic therapy, including a BTK inhibitor or B.) chronic lymphocytic leukemia or small lymphocytic lymphoma who have received at least 2 prior lines of therapy, including a Bruton tyrosine kinase inhibitor and a B-cell lymphoma 2 inhibitor.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **KALYDECO**

#### **Products Affected**

KALYDECO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **KERENDIA**

#### **Products Affected**

• KERENDIA

PA Criteria	Criteria Details
Exclusion Criteria	none
Required Medical Information	Diagnosis of Chronic kidney disease associated with Type 2 diabetes mellitus AND patient is currently receiving the following standard of care A.) A maximally tolerated dose of ACE inhibitor, ARB, or a combination medication containing an ACE or ARB AND B.) Antidiabetic agent (e.g., metformin or an agent containing metformin, SGLT2 inhibitor, GLP-1 RA)
Age Restrictions	none
Prescriber Restrictions	none
Coverage Duration	12 months
Other Criteria	none
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **KISQALI**

#### **Products Affected**

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)

• KISQALI (600 MG DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# KISQALI FEMARA

#### **Products Affected**

- KISQALI FEMARA (200 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISOALI FEMARA (400 MG DOSE)

KISQALI FEMARA (400 MG DOSE)	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following A.) The patient is pre-or perimenopausal woman or male and the requested drug will be used as initial endocrine-based therapy, B.) The patient is postmenopausal, the requested drug will be used as initial endocrine-based therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **KORLYM**

#### **Products Affected**

• KORLYM

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, B.) Concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, C.) History of unexplained vaginal bleeding, D.) Endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and must meet all of the following 1.) Used to control hyperglycemia secondary to hypercortisolism, AND 2.) Patient has failed or is not a candidate for surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **KOSELUGO**

#### **Products Affected**

KOSELUGO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of neurofibromatosis type 1 (NF1) in a patient who has symptomatic, inoperable plexiform neurofibromas (PN)
Age Restrictions	2 years of age to 17 years of age
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **KRAZATI**

#### **Products Affected**

KRAZATI

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **LAPATINIB**

#### **Products Affected**

• lapatinib ditosylate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND meets one of the following A.) Used in combination with capecitabine in a patient who has received prior therapy including an anthracycline, a taxane, and trastuzumab, OR B.) Used in combination with letrozole in a postmenopausal female for whom hormonal therapy is indicated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **LENALIDOMIDE**

#### **Products Affected**

• lenalidomide

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **LENVIMA**

#### **Products Affected**

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma, in combination with everolimus, following one prior anti-angiogenic therapy, C.) Unresectable hepatocellular carcinoma, first-line therapy, D.) Advanced endometrial carcinoma that is not microsatellite instability-high or mismatch repair deficient, in combination with pembrolizumab, when disease has progressed following prior systemic therapy and patient is not a candidate for curative surgery or radiation, or E.) Advanced renal cell carcinoma, in combination with pembrolizumab and used as first-line therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **LEUPROLIDE**

#### **Products Affected**

- ELIGARD SUBCUTANEOUS KIT 22.5 MG, 30 MG, 45 MG, 7.5 MG
- leuprolide acetate
- LEUPROLIDE ACETATE (3 MONTH)
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet one of the following: 1.) Eligard only: Advanced or metastatic prostate cancer, or 2.) For Lupron depot and leuprolide products only: A.) Advanced or metastatic prostate cancer and patient has failed or is intolerant to Eligard or Trelstar (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, and 45 mg 6-month depots only), B.) Endometriosis (3.75 mg 1-month and 11.25 mg 3-month depots only), C.) Anemia due to uterine leiomyomata (Fibroids) (3.75 mg 1-month and 11.25 mg 3-month depots only) and patient is preoperative, or D.) Central precocious puberty (idiopathic or neurogenic) in children
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

# **L-GLUTAMINE**

#### **Products Affected**

• l-glutamine

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# LIDOCAINE EXT

#### **Products Affected**

• lidocaine external ointment

• lidocaine hcl

PA Criteria	Criteria Details
Exclusion Criteria	Amide hypersensitivity
Required Medical Information	The requested drug will be used for or topical anesthesia of skin or mucous membranes
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **LINEZOLID**

#### **Products Affected**

• linezolid intravenous

• linezolid oral tablet

• linezolid oral suspension reconstituted

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI
Required Medical Information	Diagnosis of one of the following A.) Community acquired pneumonia, B.) Hospital-acquired pneumonia, C.) Vancomycin-resistant Enterococcus faecium infection, D.) Complicated skin and skin structure infections, or E.) Uncomplicated skin and skin structure infections
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	1 month
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **LIVTENCITY**

#### **Products Affected**

• LIVTENCITY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of post-transplant cytomegalovirus (CMV) infection/disease that is refractory to treatment (with or without genotypic resistance) with ganciclovir, valganciclovir, cidofovir or foscarnet
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **LONSURF**

### **Products Affected**

LONSURF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, or B.) Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **LORBRENA**

### **Products Affected**

• LORBRENA ORAL TABLET 100 MG, 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A4 inducers
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **LUMAKRAS**

### **Products Affected**

• LUMAKRAS ORAL TABLET 120 MG, 320 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC) as determined by an FDA-approved test, and patient has received at least one prior systemic therapy (e.g., immune checkpoint inhibitor, platinum-based chemotherapy)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **LYNPARZA**

### **Products Affected**

LYNPARZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) HER2-negative, deleterious or suspected deleterious germline BRCA mutated high-risk early or metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), C.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, D.) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen, E.) Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA-mutation and/or genomic instability AND are using in combination with bevacizumab for maintenance treatment, F.) Deleterious or suspected deleterious germline or somatic homologous recombination repair gene mutated metastatic castration-resistant prostate cancer in patients who have progressed following prior treatment with enzalutamide or abiraterone, or G.) Deleterious or suspected deleterious BRCA-mutated metastatic castration-resistant prostate cancer in combination with abiraterone and prednisone or prednisolone
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **LYTGOBI**

#### **Products Affected**

- LYTGOBI (12 MG DAILY DOSE)
- LYTGOBI (16 MG DAILY DOSE)

• LYTGOBI (20 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of unresectable, locally advanced or metastatic intrahepatic cholangiocarcinoma harboring fibroblast growth factor receptor 2 (FGFR2) gene fusions or other rearrangements and previously treated
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **MATULANE**

### **Products Affected**

MATULANE

PA Criteria	Criteria Details
Exclusion Criteria	Inadequate marrow reserve
Required Medical Information	Diagnosis of Hodgkin's Disease, Stages III and IV and used in combination with other anticancer drugs
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **MAYZENT**

#### **Products Affected**

• MAYZENT ORAL TABLET 0.25 MG, 1 • MAYZENT STARTER PACK MG, 2 MG

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) CYP2C9*3/*3 genotype, B.) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, Class III-IV heart failure, or C.) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **MEKINIST**

### **Products Affected**

MEKINIST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy, D.) Metastatic non-small cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib, E.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with dabrafenib, and have progressed following prior treatment and have no satisfactory alternative treatment options, F.) Low-grade glioma with a BRAF V600E mutation and require systemic therapy, in combination with dabrafenib
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **MEKTOVI**

### **Products Affected**

MEKTOVI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic malignant melanoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib or B.) Metastatic non-small cell lung cancer with a BRAF V600E mutation as detected by an FDA-approved test AND used in combination with encorafenib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **MIFEPRISTONE**

### **Products Affected**

• mifepristone

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, B.) Concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, C.) History of unexplained vaginal bleeding, D.) Endometrial hyperplasia with atypia or endometrial carcinoma
Required Medical Information	Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and must meet all of the following 1.) Used to control hyperglycemia secondary to hypercortisolism, and 2.) Patient has failed or is not a candidate for surgery
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **MIGLUSTAT**

### **Products Affected**

• miglustat

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **NAYZILAM**

### **Products Affected**

NAYZILAM

PA Criteria	Criteria Details
Exclusion Criteria	Acute narrow angle glaucoma
Required Medical Information	Diagnosis of epilepsy and documentation of acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e. seizure clusters, acute repetitive seizures), that are distinct from a patient's usual seizure pattern
Age Restrictions	12 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	This criteria applies to new starts only
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **NERLYNX**

### **Products Affected**

NERLYNX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Early stage HER2-positive breast cancer and used following adjuvant trastuzumab therapy, or B.) Advanced or metastatic HER2-positive breast cancer, used in combination with capecitabine, AND patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **NINLARO**

### **Products Affected**

NINLARO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of multiple myeloma, used in combination with lenalidomide and dexamethasone, AND patient has history of at least 1 prior therapy. Examples include bortezomib, cyclophosphamide, Kyprolis (carfilzomib intravenous infusion), lenalidomide, Darzalex (daratumumab intravenous infusion).
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **NUBEQA**

### **Products Affected**

NUBEQA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Non-metastatic, castration-resistant prostate cancer (nmCRPC) or B.) Metastatic hormone-sensitive prostate cancer in combination with docetaxel. For treatment of nmCRPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **NUEDEXTA**

### **Products Affected**

NUEDEXTA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy, F.) History of quinine-, mefloquine-, or quinidine-induced thrombocytopenia, bone marrow depression, or lupus-like syndrome
Required Medical Information	Diagnosis of pseudobulbar affect
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **NUPLAZID**

### **Products Affected**

NUPLAZID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Parkinson's disease and both of the following apply A.) Used for treatment of hallucinations and/or delusions associated with Parkinson's disease psychosis, and B.) Diagnosis of Parkinson's disease was made prior to the onset of psychotic symptoms
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OCTREOTIDE**

### **Products Affected**

• octreotide acetate

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **ODOMZO**

### **Products Affected**

ODOMZO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OFEV**

### **Products Affected**

OFEV

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OGSIVEO**

#### **Products Affected**

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of progressing desmoid tumors who require systemic treatment
Age Restrictions	18 years or older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OJEMDA**

#### **Products Affected**

- OJEMDA ORAL SUSPENSION RECONSTITUTED
- OJEMDA ORAL TABLET

RECONSTITUTED	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory pediatric low-grade glioma harboring a BRAF fusion or rearrangement, or BRAF V600 mutation
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OJJAARA**

### **Products Affected**

OJJAARA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate or high-risk myelofibrosis (MF), including primary MF or secondary MF [postpolycythemia vera (PV) and postessential thrombocythemia (ET)], in adults with anemia.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ONUREG**

### **Products Affected**

• ONUREG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acute myeloid leukemia (AML) used in maintenance treatment for adult patients who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ORENCIA**

### **Products Affected**

ORENCIA

### • ORENCIA CLICKJECT

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, or D.) Prophylaxis of acute graft versus host disease (aGVHD), in combination with a calcineurin inhibitor and methotrexate, in patients undergoing hematopoietic stem cell transplantation (HSCT) from a matched or 1 allele-mismatched unrelated-donor.
Age Restrictions	For RA: 18 years of age or older. All other indications 2 years of age or older.
Prescriber Restrictions	For RA and PJIA: Rheumatologist, For PsA: Dermatologist or Rheumatologist, For aGVHD: Hematologist or Transplant Specialist.
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ORGOVYX**

### **Products Affected**

ORGOVYX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ORKAMBI**

### **Products Affected**

ORKAMBI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ORSERDU**

#### **Products Affected**

• ORSERDU ORAL TABLET 345 MG, 86 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced or metastatic, ER-positive, HER2-negative, ESR1-mutated, breast cancer in postmenopausal women or adult man after at least 1 line of endocrine therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **OTEZLA**

### **Products Affected**

- OTEZLA ORAL TABLET
- OTEZLA ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **PANRETIN**

### **Products Affected**

• PANRETIN

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **PAZOPANIB**

### **Products Affected**

• pazopanib hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma (RCC), or B.) Advanced soft tissue sarcoma (STS), has received prior chemotherapy and does not have adipocytic soft tissue sarcoma (STS) or gastrointestinal stromal tumors (GIST)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **PEGYLATED INTERFERON**

### **Products Affected**

• PEGASYS

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

### **PEMAZYRE**

### **Products Affected**

PEMAZYRE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with confirmed fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test or B.) Relapsed or refractory myeloid/lymphoid neoplasms with fibroblast growth factor receptor 1 rearrangement
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, gastroenterologist, or hepatologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **PENICILLAMINE**

### **Products Affected**

• penicillamine

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Breastfeeding, B.) Hypersensitivity to penicillamine products, C.) Penicillamine-related aplastic anemia/agranulocytosis, or D.) Rheumatoid arthritis patients with history or evidence of renal insufficiency
Required Medical Information	Diagnosis of one of the following A.) Cystinuria, B.) Severe, active rheumatoid arthritis, or C.) Wilson's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **PIQRAY**

#### **Products Affected**

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)

• PIQRAY (300 MG DAILY DOSE)

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **PIRFENIDONE**

#### **Products Affected**

• pirfenidone

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of idiopathic pulmonary fibrosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **POMALYST**

#### **Products Affected**

POMALYST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) AIDS-related Kaposi sarcoma and patient has failure on highly active antiretroviral therapy (HAART), B.) Kaposi sarcoma in HIV-negative adults, or C.) Multiple myeloma and in combination with dexamethasone in adults who have received at least 2 prior therapies (including lenalidomide and a proteasome inhibitor) and have demonstrated disease progression on or within 60 days of completion of the last therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **POSACONAZOLE**

#### **Products Affected**

posaconazole

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **PREVYMIS**

#### **Products Affected**

• PREVYMIS

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use with pimozide or ergot alkaloids (ergotamine, dihydroergotamine), B.) Concomitant use with pitavastatin or simvastatin when coadministered with cyclosporine
Required Medical Information	Diagnosis of one of the following A.) Prophylaxis of cytomegalovirus (CMV) infection and disease in adult CMV-seropositive recipients [R+] of an allogeneic hematopoietic stem cell transplant, or B.) Prophylaxis of CMV disease in adult kidney transplant recipients at high risk (Donor CMV seropositive/Recipient CMV seronegative [D+/R-])
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	6 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **PROLIA**

#### **Products Affected**

PROLIA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Must meet all of the following A.) Diagnosis of one of the following 1.) Treatment of osteoporosis in postmenopausal females at high risk of fracture, 2.) Treatment of osteoporosis in males at high risk of fracture 3.) Treatment of bone loss in males receiving androgen-deprivation therapy for nonmetastatic prostate cancer, 4.) Treatment of bone loss in females receiving aromatase inhibitor therapy for breast cancer, or 5.) Treatment of glucocorticoid-induced osteoporosis in patients at high risk of fracture who are initiating or continuing systemic glucocorticoids at a daily dose greater than or equal to 7.5 mg of prednisone for an anticipated duration of at least 6 months B.) Bone density scan (DEXA) documenting a T-score less than 1.0 at lumbar spine, total hip, femoral neck, or 33% radius and member has high risk factors (high risk defined as osteoporotic fracture history, multiple risk factors for fracture, or failure of or intolerance to other available osteoporosis therapy) or T-score less than or equal to -2.5 in the lumbar spine, total hip, femoral neck, or 33% radius
Age Restrictions	none
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

### **PROMACTA**

#### **Products Affected**

• PROMACTA ORAL PACKET 12.5 MG, • PROMACTA ORAL TABLET 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy or in combination with standard immunosuppressive therapy
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **PULMOZYME**

#### **Products Affected**

• PULMOZYME

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **PURIXAN**

#### **Products Affected**

• PURIXAN

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of acute lymphocytic leukemia, 2.) Using in combination with methotrexate, 3.) Patient has tried/failed or has contraindication/intolerance to mercaptopurine tablets
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **PYRIMETHAMINE**

#### **Products Affected**

• pyrimethamine

PA Criteria	Criteria Details
Exclusion Criteria	Documented megaloblastic anemia due to folate deficiency
Required Medical Information	Diagnosis of Toxoplasmosis and treatment in combination with a sulfonamide
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	10 weeks
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **QINLOCK**

#### **Products Affected**

• QINLOCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced gastrointestinal stromal tumor (GIST) and patient has received prior treatment with 3 or more kinase inhibitors, including imatinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **QUININE SULFATE**

#### **Products Affected**

• quinine sulfate

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **RAVICTI**

#### **Products Affected**

• RAVICTI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of urea cycle disorders
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **REGRANEX**

### **Products Affected**

REGRANEX

PA Criteria	Criteria Details
Exclusion Criteria	Known neoplasm at the site of application
Required Medical Information	Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply
Age Restrictions	16 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **REPATHA**

#### **Products Affected**

• REPATHA

- REPATHA SURECLICK
- REPATHA PUSHTRONEX SYSTEM

• KEI ATTIA I USITIKONEA STSTEM	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH), B.) homozygous familial hypercholesterolemia, C.) established cardiovascular disease and myocardial infarction prophylaxis, stroke prophylaxis, or coronary revascularization prophylaxis is required, or D.) clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following 1.) acute coronary syndrome, 2.) history of myocardial infarction, 3.) stable/unstable angina, 4.) coronary or other arterial revascularization, 5.) stroke, 6.) transient ischemic stroke (TIA), or 7.) peripheral arterial disease presumed to be atherosclerotic region
Age Restrictions	10 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

### **RETEVMO**

#### **Products Affected**

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy, B.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC), C.) Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and are refractory to radioactive iodine, if appropriate, or D.) Locally advanced or metastatic solid tumors with a RET gene fusion that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

# **REVLIMID**

#### **Products Affected**

• REVLIMID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

### **REZLIDHIA**

#### **Products Affected**

• REZLIDHIA

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **REZUROCK**

#### **Products Affected**

REZUROCK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of chronic graft-vs-host disease and patient has failed at least 2 prior lines of systemic therapy (e.g., prednisone, methotrexate, cyclosporine, tacrolimus, mycophenolate, Imbruvica (ibrutinib), Jakafi (ruxolitinib), etc.)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **RILUZOLE**

#### **Products Affected**

• riluzole

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of amyotrophic lateral sclerosis (ALS)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **RINVOQ**

#### **Products Affected**

• RINVOQ

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **RINVOQ LQ**

#### **Products Affected**

• RINVOQ LQ

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Active psoriatic arthritis with inadequate response or intolerance to 1 or more TNF blockers, or B.) Active polyarticular juvenile idiopathic arthritis with inadequate response or intolerance to 1 or more TNF blockers
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a rheumatologist
Coverage Duration	Plan year
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ROZLYTREK**

#### **Products Affected**

- ROZLYTREK ORAL CAPSULE 100 MG, 200 MG
- ROZLYTREK ORAL PACKET

WIG, 200 WIG	
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) ROS1-positive metastatic non-small cell lung cancer (NSCLC), or B.) Solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity, and have either progressed following treatment or have no satisfactory alternative therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	Verify the pregnancy status of females of reproductive potential prior to initiating ROZLYTREK
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **RUBRACA**

#### **Products Affected**

• RUBRACA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, used as maintenance treatment, and patient is in complete or partial response to platinum-based chemotherapy, or B.) Deleterious BRCA mutation (germline and/or somatic)-associated metastatic castration-resistant prostate cancer and patient has been treated with androgen receptor-directed therapy and a taxane-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **RUFINAMIDE**

#### **Products Affected**

• rufinamide oral suspension

PA Criteria	Criteria Details
Exclusion Criteria	Familial Short QT Syndrome
Required Medical Information	Diagnosis of seizures associated with Lennox-Gastaut syndrome and used as adjunctive treatment
Age Restrictions	1 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **RYDAPT**

#### **Products Affected**

• RYDAPT

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **SAPROPTERIN**

#### **Products Affected**

• sapropterin dihydrochloride

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Initial: 2 months, Renewal: Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SCEMBLIX**

#### **Products Affected**

• SCEMBLIX

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs), or B.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) with the T315I mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SIGNIFOR**

#### **Products Affected**

• SIGNIFOR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Cushing disease and patient has had inadequate response to or is not a candidate for surgery. For renewal: Documentation of a clinically meaningful reduction in 24-hour urinary free cortisol (UFC) levels or improvement in signs or symptoms of the disease
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **SILDENAFIL**

#### **Products Affected**

• sildenafil citrate oral tablet 20 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Nitrate therapy, including intermittent use, B.) Concomitant use with riocguat or other guanylate cyclase stimulators, C.) Concomitant use with HIV protease inhibitors or elvitegravir/cobicistat/tenofovir/emtricitabine
Required Medical Information	Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist or pulmonologist
Coverage Duration	12 months
Other Criteria	Continuation of therapy: patient is tolerating and responding to medication and has improved exercise capacity or a delay in clinical worsening.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SIRTURO**

#### **Products Affected**

• SIRTURO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Must meet all of the following 1.) Diagnosis of pulmonary multidrug resistant tuberculosis (MDR-TB) and 2.) Used in combination with at least 3 other antibiotics for the treatment of pulmonary multi-drug resistant tuberculosis
Age Restrictions	5 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	24 weeks
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **SKYRIZI**

#### **Products Affected**

- SKYRIZI (150 MG DOSE)
- SKYRIZI PEN
- SKYRIZI SUBCUTANEOUS SOLUTION CARTRIDGE 180 MG/1.2ML, 360 MG/2.4ML
- SKYRIZI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **SODIUM OXYBATE**

#### **Products Affected**

• sodium oxybate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency
Required Medical Information	Diagnosis of one of the following A.) Narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to a central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, methylphenidate) or a CNS wakefulness promoting drug (e.g., armodafinil, modafinil), or B.) Cataplexy and narcolepsy
Age Restrictions	7 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SOLTAMOX**

#### **Products Affected**

SOLTAMOX

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant coumarin-type anticoagulant therapy, B.) history of thromboembolic disease such as DVT or PE
Required Medical Information	Diagnosis of breast cancer and documentation of inability to swallow tablet formulation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **SOMAVERT**

#### **Products Affected**

• SOMAVERT SUBCUTANEOUS SOLUTION RECONSTITUTED 10 MG, 15 MG, 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of acromegaly confirmed by high pretreatment insulin-like growth factor-1 (IGF-1) level for age and/or gender based on the laboratory reference range and patient has had an inadequate response to or is ineligible for surgery or radiation therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an endocrinologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SORAFENIB**

#### **Products Affected**

• sorafenib tosylate

PA Criteria	Criteria Details
Exclusion Criteria	Squamous cell lung cancer being treated with carboplatin and paclitaxel
Required Medical Information	Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Unresectable hepatocellular carcinoma
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **SPRYCEL**

#### **Products Affected**

 SPRYCEL ORAL TABLET 100 MG, 140 MG, 20 MG, 50 MG, 70 MG, 80 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, B.) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy, C.) Ph+ acute lymphoblastic leukemia (ALL) with resistance or intolerance to prior therapy, or D.) Newly diagnosed Ph+ ALL in combination with chemotherapy
Age Restrictions	1 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **STELARA**

#### **Products Affected**

- STELARA SUBCUTANEOUS SOLUTION
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 45 MG/0.5ML, 90 MG/ML

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severely active Crohn's disease, B.) Moderate to severe plaque psoriasis, C.) Active psoriatic arthritis, or D.) Moderate to severe active ulcerative colitis. For moderate to severe plaque psoriasis: 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected)
Age Restrictions	For PSO or PsA: 6 years of age or older, For all other indications: 18 years of age or older
Prescriber Restrictions	For PsO: Dermatologist, For PsA: Dermatologist or Rheumatologist, For CD and UC: Gastroenterologist.
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

PA Criteria	Criteria Details
Off-Label Uses	N/A
Part B Prerequisite	No

## **STIVARGA**

#### **Products Affected**

STIVARGA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic colorectal cancer in patients previously treated with fluoropyrimidine, oxaliplatin, and irinotecan containing chemotherapy, anti-VEGF therapy, and if RAS wild type, anti-EGFR therapy, B.) Liver carcinoma in patients previously treated with sorafenib, or C.) Locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with imatinib and sunitinib
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SUNITINIB**

#### **Products Affected**

• sunitinib malate

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) Pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease, C.) Advanced renal cell carcinoma, or D.) Renal cell carcinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **SYMLIN**

### **Products Affected**

• SYMLINPEN 120

#### • SYMLINPEN 60

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Confirmed diagnosis of gastroparesis, B.) Hypoglycemia unawareness
Required Medical Information	Diagnosis of type 1 or type 2 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **SYNAREL**

#### **Products Affected**

• SYNAREL

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Breastfeeding, or B.) undiagnosed abnormal vaginal bleeding
Required Medical Information	Diagnosis of one of the following A.) Central precocious puberty, or B.) Endometriosis
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TABRECTA**

### **Products Affected**

TABRECTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TADALAFIL**

#### **Products Affected**

• tadalafil

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) concurrent use of nitrates, including intermittent use, or B.) diagnosis of erectile dysfunction without signs and symptoms of BPH
Required Medical Information	Diagnosis of benign prostatic hyperplasia (BPH) and patient has experienced intolerance to or treatment failure to ONE alpha-blocker (e.g., doxazosin, prazosin, tamsulosin) AND to ONE 5-alpha reductase inhibitor (e.g., dutasteride, finasteride).
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a urologist
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TAFINLAR**

### **Products Affected**

TAFINLAR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options, B.) Metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation, D.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with trametinib, and have progressed following prior treatment and have no satisfactory alternative treatment options, or E.) Low-grade glioma with a BRAF V600E mutation and require systemic therapy, in combination with trametinib
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

# **TAGRISSO**

#### **Products Affected**

TAGRISSO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, B.) Metastatic non-small cell lung cancer with T790M EGFR mutation (as confirmed by an FDA-approved test) AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor therapy, C.) Non-small cell lung cancer (NSCLC) with tumor epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations (as confirmed by an FDA-approved test) AND patient requires adjuvant therapy after tumor resection, or D.) First-line treatment of adult patients with locally advanced or metastatic non-small cell lung cancer whose tumors have epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations, as detected by an FDA-approved test, in combination with pemetrexed and platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

### **TAKHZYRO**

#### **Products Affected**

- TAKHZYRO SUBCUTANEOUS SOLUTION
- TAKHZYRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 150 MG/ML, 300 MG/2ML

PA Criteria	Criteria Details
1 A CITICITA	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following and used as routine prophylaxis A.) Hereditary angioedema (HAE) with C1 inhibitor deficiency (Type 1) confirmed by laboratory testing, B.) HAE with C1 inhibitor dysfunction (Type 2) confirmed by laboratory testing, or C.) HAE with normal C1 inhibitor (Type 3) confirmed by laboratory testing and one of the following 1.) Positive test for an F12, angiopoietin-1, or plasminogen gene mutation, or 2.) Family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a hematologist, immunologist, or allergist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TALZENNA**

#### **Products Affected**

TALZENNA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm), human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer, or B.) Homologous recombination repair gene-mutated metastatic castration-resistant prostate cancer in combination with enzulatamide
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TASIGNA**

#### **Products Affected**

TASIGNA

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of one of the following A.) Newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (CML), B.) Chronic phase or accelerated phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior therapy that included imatinib, or C.) Chronic phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior tyrosine-kinase inhibitor therapy
Age Restrictions	1 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TAVNEOS**

### **Products Affected**

TAVNEOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe active anti-neutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis (granulomatosis with polyangiitis [GPA] and microscopic polyangiitis [MPA]) and both of the following apply 1.) Used as adjunctive treatment, and 2.) Used in combination with standard therapy including glucocorticoids
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TAZAROTENE**

#### **Products Affected**

• tazarotene external cream

TAZORAC

• tazarotene external gel

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Acne vulgaris and patient has trial with at least one generic topical acne product, or B.) Stable moderate to severe plaque psoriasis with 20% or less body surface area involvement and patient has trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs)
Age Restrictions	12 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TAZVERIK**

### **Products Affected**

TAZVERIK

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection, B.) Relapsed or refractory follicular lymphoma in patients whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies, or C.) Relapsed or refractory follicular lymphoma in patients who have no satisfactory alternative treatment options
Age Restrictions	16 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TEPMETKO**

#### **Products Affected**

TEPMETKO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer (NSCLC) with mesenchymal-epithelial transition (MET) exon 14 skipping alterations
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **TERIPARATIDE**

#### **Products Affected**

• TERIPARATIDE (RECOMBINANT)

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

## **TETRABENAZINE**

#### **Products Affected**

• tetrabenazine oral tablet 12.5 mg, 25 mg

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine
Required Medical Information	Diagnosis of chorea associated with Huntington's disease
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **THALOMID**

#### **Products Affected**

• THALOMID

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Multiple myeloma that is newly diagnosed, or B.) Erythema nodosum leprosum (ENL)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, infectious disease specialist, or dermatologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TIBSOVO**

#### **Products Affected**

• TIBSOVO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory acute myeloid leukemia with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test), B.) Previously treated, locally advanced or metastatic cholangiocarcinoma with an isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test.), C.) Acute myeloid leukemia (newly-diagnosed) with susceptible isocitrate dehydrogenase-1 mutation and meets one of the following: 1.) Patient is 75 years of age or older, or 2.) Patient has comorbidities that preclude intensive induction chemotherapy, or D.) Relapsed or refractory myelodysplastic syndromes with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a gastroenterologist, hematologist, hepatologist, or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

# **TOBI**

### **Products Affected**

• TOBI PODHALER

PA Criteria	Criteria Details
Exclusion Criteria	Known sensitivity to any aminoglycoside
Required Medical Information	Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has suspected or confirmed Pseudomonas aeruginosa infection in the lungs
Age Restrictions	6 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TOPICAL RETINOIDS**

#### **Products Affected**

• adapalene

• tretinoin external

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of mild to moderate acne vulgaris
Age Restrictions	PA applies to patients older than 26 years of age
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### TOPICAL TESTOSTERONE

#### **Products Affected**

- testosterone transdermal gel 10 mg/act (2%), 12.5 mg/act (1%), 20.25 mg/1.25gm (1.62%), 20.25 mg/act (1.62%), 25
- mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%), 50 mg/5gm (1%)
- testosterone transdermal solution

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Carcinoma of the breast (males only), or B.) Known or suspected carcinoma of the prostate
Required Medical Information	Diagnosis of one of the following A.) Hypogonadotropic hypogonadism, or B.) Primary hypogonadism. Diagnosis of hypogonadism must be confirmed by a low-for-age serum testosterone (total or free) level defined by the normal laboratory reference value
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TOREMIFENE**

#### **Products Affected**

• toremifene citrate

PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Acquired or congenital long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia
Required Medical Information	Diagnosis of metastatic breast cancer and patient must have previous inadequate response or intolerance to tamoxifen
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	6 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TRELSTAR**

#### **Products Affected**

• TRELSTAR MIXJECT INTRAMUSCULAR SUSPENSION

RECONSTITUTED 11.25 MG, 22.5 MG, 3.75 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of advanced prostate cancer
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TREMFYA**

### **Products Affected**

• TREMFYA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe plaque psoriasis, B.) Active psoriatic arthritis. For moderate to severe plaque psoriasis: 1) at least 3% of body surface area (BSA) is affected OR crucial body areas (e.g., feet, hands, face, neck, groin, intertriginous areas) are affected at the time of diagnosis, AND 2) the patient meets any of the following: a) the patient has experienced an inadequate treatment response or intolerance to either phototherapy (e.g., UVB, PUVA) or pharmacologic treatment with methotrexate, cyclosporine, or acitretin, b) pharmacologic treatment with methotrexate, cyclosporine, or acitretin is contraindicated, c) the patient has severe psoriasis that warrants a biologic as first-line therapy (i.e., at least 10% of the BSA or crucial body areas [e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas] are affected).
Age Restrictions	18 years of age or older
Prescriber Restrictions	For PsO: Dermatologist, For PsA Dermatologist or Rheumatologist
Coverage Duration	Plan Year
Other Criteria	Only for biologic therapy-naive patients: Physician's certification stating tuberculosis (TB) negative status or negative result of a TB detection test with a reading date less than 12 months before the start of therapy or chest x ray.
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

MCS Classicare Formulary 2 - Formulary ID: 25511 Effective date: 01/01/2025 Last Updated: 09/26/2024

## **TRIENTINE**

#### **Products Affected**

• trientine hcl

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Wilson's disease in patients that are intolerant to penicillamine
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TRIKAFTA**

#### **Products Affected**

TRIKAFTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of cystic fibrosis (CF) and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene or a mutation in the CFTR gene that is responsive to elexacaftor/tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test
Age Restrictions	2 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TRUQAP**

#### **Products Affected**

• TRUQAP

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, locally advanced or metastatic breast cancer with 1 or more PIK3CA/AKT1/PTEN-alterations as detected by an FDA-approved test and, A.) patient has had disease progression following 1 or more endocrine-based regimen(s) in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy, and B.) will be used in combination with fulvestrant injection.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **TUKYSA**

#### **Products Affected**

TUKYSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following: A.) advanced unresectable or metastatic HER2-positive breast cancer (including brain metastases) in patients who have received one or more prior anti-HER2-based regimens in the metastatic setting and drug is being used in combination with trastuzumab and capecitabine, or B.) unresectable or metastatic RAS wild-type, HER2-positive colorectal cancer that has progressed following treatment with fluoropyrimidine, oxaliplatin, and irinotecan-based chemotherapy and drug is being used in combination with trastuzumab
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TURALIO**

### **Products Affected**

• TURALIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery. Liver function tests prior to initiation.
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **TYMLOS**

#### **Products Affected**

• TYMLOS

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of osteoporosis in men or postmenopausal women and one of the following A.) osteoporotic fracture or multiple risk factors for fracture, or B.) previous trial of/or contraindication to bisphosphonate
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan year (Maximum 24 month treatment per patient lifetime)
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **UBRELVY**

#### **Products Affected**

UBRELVY

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin)
Required Medical Information	Diagnosis of migraine disorder with or without aura and patient has documented trial, inadequate response, or contraindication to at least 1 generic formulary triptan
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **VALCHLOR**

#### **Products Affected**

VALCHLOR

PA Criteria	Criteria Details
Exclusion Criteria	PA Criteria: Pending CMS Approval
Required Medical Information	PA Criteria: Pending CMS Approval
Age Restrictions	PA Criteria: Pending CMS Approval
Prescriber Restrictions	PA Criteria: Pending CMS Approval
Coverage Duration	PA Criteria: Pending CMS Approval
Other Criteria	PA Criteria: Pending CMS Approval
Indications	PA Criteria: Pending CMS Approval
Off-Label Uses	PA Criteria: Pending CMS Approval
Part B Prerequisite	No

# **VALTOCO**

#### **Products Affected**

• VALTOCO 20 MG DOSE

PA Criteria	Criteria Details
Exclusion Criteria	Acute narrow angle glaucoma
Required Medical Information	Diagnosis of epilepsy and documentation of use for acute treatment of intermittent, stereotypic episodes of frequent seizure activity (i.e., seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern.
Age Restrictions	6 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VANFLYTA**

### **Products Affected**

VANFLYTA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Patient must have all of the following A.) Newly diagnosed acute myeloid leukemia with FLT3-ITD mutation, B.) Used in combination with standard cytarabine and anthracycline induction and cytarabine consolidation, and as maintenance monotherapy following consolidation chemotherapy, and C.) Must be enrolled in the VANFLYTA REMS program
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VENCLEXTA**

#### **Products Affected**

VENCLEXTA

#### • VENCLEXTA STARTING PACK

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL
Required Medical Information	Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL), or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VEOZAH**

### **Products Affected**

VEOZAH

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of moderate to severe menopausal vasomotor symptoms (VMS) and A.) Initial treatment: The patient is experiencing 7 or more hot flashes per day and has a trial, failure or contraindication to hormonal therapy such as estradiol patch or oral conjugated estrogens, or B.) Continuing therapy: The patient has continued need for VMS treatment and experienced a reduction in VMS frequency or severity due to Veozah treatment
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **VERQUVO**

#### **Products Affected**

VERQUVO

PA Criteria	Criteria Details
Exclusion Criteria	Concomitant use of other soluble guanylate cyclase (sGC) stimulators
Required Medical Information	Diagnosis of chronic heart failure (HF), NYHA Class II to IV and all of the following 1.) Left ventricular ejection fraction less than 45%, 2.) Previous hospitalization for HF within 6 months or outpatient IV diuretic treatment for HF within 3 months
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a cardiologist
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VERZENIO**

### **Products Affected**

VERZENIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, node-positive, early breast cancer and ALL of the following: 1.) Patient is at high risk of recurrence, and 2.) Requested drug will be used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) for adjuvant treatment, OR B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following 1.) Used in combination with fulvestrant in a patient with disease progression following endocrine therapy, 2.) Used as monotherapy in a patient with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting, or 3.) For postmenopausal women, and men, used as initial endocrine-based treatment in combination with an aromatase inhibitor
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **VIGABATRIN**

#### **Products Affected**

- vigabatrin
- vigadrone

vigpoder

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Infantile spasms, or B.) Refractory complex partial seizures and the drug is being used as adjunctive therapy in patients who have responded inadequately to two alternative treatments
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **VIJOICE**

### **Products Affected**

VIJOICE

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of severe manifestations of PIK3CA-Related Overgrowth Spectrum (PROS) in patients who require systemic therapy
Age Restrictions	2 years of age and older
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VITRAKVI**

#### **Products Affected**

• VITRAKVI ORAL CAPSULE 100 MG, • VITRAKVI ORAL SOLUTION 25 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion positive solid tumors without a known acquired resistance mutation and used in patients with unsatisfactory alternative treatments or who have progressed following treatment
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VIZIMPRO**

### **Products Affected**

VIZIMPRO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **VONJO**

### **Products Affected**

• VONJO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of intermediate or high-risk primary or secondary myelofibrosis in adults AND a platelet count less than 50 X 10(9)/L
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **VORICONAZOLE**

#### **Products Affected**

- VORICONAZOLE INTRAVENOUS
- voriconazole oral tablet 200 mg, 50 mg
- voriconazole oral suspension reconstituted

• vortconazote orai suspension reconstitutea	
PA Criteria	Criteria Details
Exclusion Criteria	Any of the following A.) Concomitant use of carbamazepine, CYP3A4 substrates (e.g., terfenadine, astemizole, cisapride, pimozide, or quinidine), B.) Concomitant use with high-dose ritonavir (400mg every 12 hours), C.) Concomitant use with ergot alkaloids, D.) Concomitant use with longacting barbiturates, E.) ConcOmitant use with rifabutin or rifampin, F.) Concomitant use with sirolimus, or G.) Concomitant use with efavirenz at standard doses of 400mg/day or higher
Required Medical Information	Diagnosis of one of the following A.) Invasive aspergillosis, B.) Candidemia, C.) Esophageal Candidiasis, D.) Invasive candidiasis of the skin and abdomen, kidney, bladder wall, and wounds, or E.) Serious fungal infection due to Scedosporium apiospermum or Fusarium species
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an infectious disease specialist
Coverage Duration	6 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

## **VOWST**

#### **Products Affected**

• VOWST

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of prevention of recurrent Clostridioides difficile infection (CDI) and one of the following 1.) Patient has completed antibiotic treatment for at least 3 CDI episodes, or 2.) Previously received Vowst and both of the follwing A.) Treatment failure (defined as the presence of CDI diarrhea within 8 weeks of first dose of Vowst and a positive stool test for C. Difficile, and B.) Patient has not received more than on treatment course of vowst which was at least 12 days and not more than 8 weeks prior
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	30 days
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **VUMERITY**

#### **Products Affected**

VUMERITY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **WELIREG**

### **Products Affected**

WELIREG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Von Hippel-Lindau (VHL) disease and therapy is required for any of the following disease associated tumors that do not require immediate surgery 1.) Renal cell carcinoma (RCC), 2.) Central nervous system (CNS) hemangioblastoma, or 3.) Pancreatic neuroendocrine tumor (pNET), or B.) Advanced renal cell carcinoma following a programmed death receptor-1 or programmed death-ligand 1 inhibitor and a vascular endothelial growth factor tyrosine kinase inhibitor
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a hematologist or oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XALKORI**

#### **Products Affected**

- XALKORI ORAL CAPSULE
- XALKORI ORAL CAPSULE SPRINKLE 150 MG, 20 MG, 50 MG

PA Criteria	Criteria Details
1 A Cilulia	Citatia Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test, B.) Relapsed or refractory systemic anaplastic large cell lymphoma that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test, or C.) Unresectable, recurrent, or refractory inflammatory myofibroblastic tumors that are anaplastic lymphoma kinase (ALK)-positive
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XCOPRI**

#### **Products Affected**

- XCOPRI (250 MG DAILY DOSE)
- XCOPRI (350 MG DAILY DOSE)
- XCOPRI ORAL TABLET 100 MG, 150 MG, 200 MG, 25 MG, 50 MG

• XCOPRI ORAL TABLET THERAPY PACK

PA Criteria	Criteria Details
Exclusion Criteria	Familial short QT syndrome
Required Medical Information	Diagnosis of partial-onset seizures
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **XDEMVY**

### **Products Affected**

• XDEMVY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Demodex blepharitis
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	12 months
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XELJANZ**

#### **Products Affected**

- XELJANZ ORAL SOLUTION
- XELJANZ XR
- XELJANZ ORAL TABLET 10 MG, 5 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis (RA), B.) Active psoriatic arthritis, C.) Moderate to severe ulcerative colitis (UC), D.) Polyarticular course juvenile idiopathic arthritis (pcJIA), or E.) Active ankylosing spondylitis. Screening for latent tuberculosis infection is required prior to initiation of treatment.
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	Screening for latent tuberculosis infection is required prior to initiation of treatment
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XERMELO**

#### **Products Affected**

XERMELO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of carcinoid syndrome diarrhea and both of the following 1.) Diarrhea is inadequately controlled by a stable dose of somatostatin analog (SSA) therapy (e.g., octreotide, lanreotide) for at least 3 months, and 2.) Used in combination with SSA therapy
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an oncologist, endocrinologist, or gastroenterologist
Coverage Duration	Plan year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XGEVA**

#### **Products Affected**

XGEVA

PA Criteria	Criteria Details
Exclusion Criteria	Hypocalcemia (calcium less than 8.0 mg/dL)
Required Medical Information	Diagnosis of one of the following A.) Bone metastases from a solid tumor, used for the prevention of skeletal related events and must have an inadequate treatment response, adverse event, or contraindication to zoledronic acid, B.) Multiple myeloma, used for the prevention of skeletal related events and must have an inadequate treatment response, adverse event, or contraindication to zoledronic acid, C.) Hypercalcemia of malignancy refractory to bisphosphonate therapy and must have an inadequate treatment response, adverse event, or contraindication to zoledronic acid, or D.) Giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	B vs D determination required per CMS guidance
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	Yes

### **XOLAIR**

#### **Products Affected**

• XOLAIR

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy for patients 12 years of age and older, B.) Moderate to severe persistent asthma in patients 6 years of age or older with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids or Dupixent, C.) Nasal polyps in patients 18 years of age or older with inadequate response to nasal corticosteroids, or D.) Reduction of allergic reactions (type I), including anaphylaxis, that may occur with accidental exposure to 1 or more foods in with IgE-mediated food allergy and is being used in conjunction with food allergen avoidance in patient 1 year of age or older
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with an allergist, dermatologist, immunologist, otolaryngologist, or pulmonologist
Coverage Duration	Initial: 6 months, Renewal: Plan Year
Other Criteria	B vs D coverage determination per CMS guidelines
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **XOSPATA**

#### **Products Affected**

XOSPATA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an Oncologist or Hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **XPOVIO**

#### **Products Affected**

- XPOVIO (100 MG ONCE WEEKLY)
- XPOVIO (40 MG ONCE WEEKLY)
- XPOVIO (40 MG TWICE WEEKLY)
- XPOVIO (60 MG ONCE WEEKLY)
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY)
- XPOVIO (80 MG TWICE WEEKLY)

Ì	(Control Williams)
PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Relapsed or refractory multiple myeloma being used in combination with dexamethasone in a patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody, B.) Multiple myeloma being used in combination with bortezomib and dexamethasone in a patient who has received at least 1 prior therapy, C.) Relapsed or refractory diffuse large B-cell lymphoma not otherwise specified, or D.) Relapsed or refractory DLBCL arising from follicular lymphoma and patient has received at least 2 lines of systemic therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

Prior Authorization Criteria

### **XTANDI**

#### **Products Affected**

XTANDI

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Castration-resistant prostate cancer (CRPC), B.) Metastatic, castration-sensitive prostate cancer (mCSPC). For treatment of CRPC and mCSPC, one of the following applies 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy, or C.) Nonmetastatic castration-sensitive prostate cancer with biochemical recurrence at high risk for metastasis
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **YONSA**

#### **Products Affected**

YONSA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of metastatic, castration-resistant prostate cancer (mCRPC) and used in combination with methylprednisolone. For treatment of mCRPC, one of the following applies: 1.) Used in combination with a gonadotropin-releasing hormone (GnRH) analog or 2) Patient has received bilateral orchiectomy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or urologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZARXIO**

#### **Products Affected**

• ZARXIO

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Chemotherapy induced febrile neutropenia (prophylaxis), B.) Severe chronic neutropenia, C.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, or D.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS)
Age Restrictions	None
Prescriber Restrictions	None
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

## **ZEJULA**

### **Products Affected**

• ZEJULA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of Advanced or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer and used as maintenance therapy in a patient who is in a complete or partial response to platinum-based chemotherapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZELBORAF**

#### **Products Affected**

ZELBORAF

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of one of the following A.) Unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test, or B.) Erdheim-Chester disease and patient has documented BRAF V600 mutation
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

# **ZOLINZA**

#### **Products Affected**

ZOLINZA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of primary cutaneous T-cell lymphoma (CTCL) in patients who have progressive, persistent or recurrent disease on or following two systemic therapies (e.g., bexarotene, romidepsin, etc)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZTALMY**

### **Products Affected**

• ZTALMY

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD)
Age Restrictions	None
Prescriber Restrictions	Prescribed by or in consultation with a neurologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZURZUVAE**

#### **Products Affected**

• ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of postpartum depression
Age Restrictions	18 years of age and older
Prescriber Restrictions	None
Coverage Duration	14 days
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZYDELIG**

#### **Products Affected**

• ZYDELIG ORAL TABLET 100 MG, 150 MG

PA Criteria	Criteria Details
Exclusion Criteria	History of toxic epidermal necrosis with any drug
Required Medical Information	Diagnosis of Chronic lymphocytic leukemia, used in combination with rituximab and patient has relapsed on at least one prior therapy
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist or hematologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### **ZYKADIA**

#### **Products Affected**

• ZYKADIA

PA Criteria	Criteria Details
Exclusion Criteria	None
Required Medical Information	Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC)
Age Restrictions	18 years of age and older
Prescriber Restrictions	Prescribed by or in consultation with an oncologist
Coverage Duration	Plan Year
Other Criteria	None
Indications	All FDA-approved Indications.
Off-Label Uses	N/A
Part B Prerequisite	No

### PART B VERSUS PART D

#### **Products Affected**

- ABILIFY ASIMTUFII INTRAMUSCULAR PREFILLED SYRINGE 720 MG/2.4ML, 960 MG/3.2ML
- ABILIFY MAINTENA INTRAMUSCULAR PREFILLED SYRINGE 300 MG, 400 MG
- ABILIFY MAINTENA INTRAMUSCULAR SUSPENSION RECONSTITUTED ER 300 MG, 400 MG
- acetylcysteine inhalation solution 10 %, 20 %
- acyclovir sodium intravenous solution 50 mg/ml
- albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml
- amphotericin b intravenous solution reconstituted 50 mg
- amphotericin b liposome intravenous suspension reconstituted 50 mg
- ampicillin sodium injection solution reconstituted 1 gm, 125 mg
- aprepitant oral capsule 125 mg, 40 mg, 80
   & 125 mg, 80 mg
- azathioprine oral tablet 50 mg
- azithromycin intravenous solution reconstituted 500 mg
- budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml
- calcitonin (salmon) nasal solution 200 unit/act
- calcitriol oral capsule 0.25 mcg, 0.5 mcg
- calcitriol oral solution 1 mcg/ml
- cefoxitin sodium intravenous solution reconstituted 1 gm, 10 gm, 2 gm
- cefuroxime sodium injection solution reconstituted 750 mg
- cefuroxime sodium intravenous solution reconstituted 1.5 gm

- cinacalcet hcl oral tablet 30 mg, 60 mg,
   90 mg
- ciprofloxacin in d5w intravenous solution 200 mg/100ml
- clindamycin phosphate injection solution 900 mg/6ml
- colistimethate sodium (cba) injection solution reconstituted 150 mg
- cromolyn sodium inhalation nebulization solution 20 mg/2ml
- CYCLOPHOSPHAMIDE ORAL CAPSULE 25 MG
- cyclophosphamide oral capsule 50 mg
- cyclophosphamide oral tablet 25 mg
- CYCLOPHOSPHAMIDE ORAL TABLET 50 MG
- cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
- cyclosporine modified oral solution 100 mg/ml
- cyclosporine oral capsule 100 mg, 25 mg
- DEXTROSE-SODIUM CHLORIDE INTRAVENOUS SOLUTION 10-0.2 %, 10-0.45 %, 2.5-0.45 %
- doxy 100 intravenous solution reconstituted 100 mg
- ENGERIX-B INJECTION SUSPENSION 20 MCG/ML
- ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML
- ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg
- fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%
- GAMMAGARD INJECTION SOLUTION 2.5 GM/25ML

Prior Authorization Criteria

- GAMMAGARD S/D LESS IGA INTRAVENOUS SOLUTION RECONSTITUTED 10 GM, 5 GM
- GAMMAPLEX INTRAVENOUS SOLUTION 10 GM/100ML, 10 GM/200ML, 20 GM/200ML, 5 GM/50ML
- GAMUNEX-C INJECTION SOLUTION 1 GM/10ML
- granisetron hcl oral tablet 1 mg
- heparin sodium (porcine) injection solution 10000 unit/ml, 20000 unit/ml, 5000 unit/ml
- HEPLISAV-B INTRAMUSCULAR SOLUTION PREFILLED SYRINGE 20 MCG/0.5ML
- IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML
- ipratropium bromide inhalation solution 0.02 %
- ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- ISOLYTE-S PH 7.4 INTRAVENOUS SOLUTION
- JYLAMVO ORAL SOLUTION 2 MG/ML
- kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%
- KCL IN DEXTROSE-NACL INTRAVENOUS SOLUTION 40-5-0.9 MEQ/L-%-%
- KCL-LACTATED RINGERS-D5W INTRAVENOUS SOLUTION 20 MEQ/L
- methotrexate sodium (pf) injection solution 50 mg/2ml
- methotrexate sodium injection solution 50 mg/2ml
- metronidazole intravenous solution 500 mg/100ml

- multiple electro type 1 ph 5.5 intravenous solution
- mycophenolate mofetil oral capsule 250 mg
- mycophenolate mofetil oral suspension reconstituted 200 mg/ml
- mycophenolate mofetil oral tablet 500 mg
- mycophenolate sodium oral tablet delayed release 180 mg, 360 mg
- ondansetron hcl oral solution 4 mg/5ml
- ondansetron hcl oral tablet 4 mg, 8 mg
- ondansetron oral tablet dispersible 4 mg, 8 mg
- paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg
- penicillin g potassium injection solution reconstituted 20000000 unit
- penicillin g sodium injection solution reconstituted 5000000 unit
- pentamidine isethionate inhalation solution reconstituted 300 mg
- pentamidine isethionate injection solution reconstituted 300 mg
- PERSERIS SUBCUTANEOUS PREFILLED SYRINGE 120 MG, 90 MG
- POTASSIUM CHLORIDE IN NACL INTRAVENOUS SOLUTION 20-0.45 MEQ/L-%, 40-0.9 MEQ/L-%
- potassium chloride in nacl intravenous solution 20-0.9 meq/l-%
- POTASSIUM CHLORIDE INTRAVENOUS SOLUTION 10 MEQ/100ML, 20 MEQ/100ML, 40 MEQ/100ML
- potassium chloride intravenous solution 2 meq/ml, 2 meq/ml (20 ml)
- potassium cl in dextrose 5% intravenous solution 20 meq/l
- PREHEVBRIO INTRAMUSCULAR SUSPENSION 10 MCG/ML
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROLASTIN-C INTRAVENOUS SOLUTION 1000 MG/20ML

- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML
- RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML
- sirolimus oral solution 1 mg/ml
- sirolimus oral tablet 0.5 mg, 1 mg, 2 mg
- tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg

- TEFLARO INTRAVENOUS SOLUTION RECONSTITUTED 400 MG, 600 MG
- TIGECYCLINE INTRAVENOUS SOLUTION RECONSTITUTED 50 MG
- tobramycin inhalation nebulization solution 300 mg/5ml
- tobramycin sulfate injection solution 10 mg/ml, 80 mg/2ml
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- ziprasidone mesylate intramuscular solution reconstituted 20 mg

#### **Details**

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

### **Alphabetical Listing**

A	amphotericin b intravenous solution
ABILIFY ASIMTUFII	reconstituted 50 mg285
INTRAMUSCULAR PREFILLED	amphotericin b liposome intravenous
SYRINGE 720 MG/2.4ML, 960	suspension reconstituted 50 mg 285
MG/3.2ML285	ampicillin sodium injection solution
ABILIFY MAINTENA	reconstituted 1 gm, 125 mg 285
INTRAMUSCULAR PREFILLED	aprepitant oral capsule 125 mg, 40 mg, 80 &
SYRINGE 300 MG, 400 MG 285	125 mg, 80 mg285
ABILIFY MAINTENA	APTIOM14
INTRAMUSCULAR SUSPENSION	ARCALYST 15
RECONSTITUTED ER 300 MG, 400	ARIKAYCE16
MG285	armodafinil42
abiraterone acetate oral tablet 250 mg, 500	AUGTYRO18
mg 1	AUSTEDO19
acetylcysteine inhalation solution 10 %, 20	AUSTEDO XR ORAL TABLET
%285	EXTENDED RELEASE 24 HOUR 12
acitretin2	MG, 18 MG, 24 MG, 30 MG, 36 MG, 42
ACTEMRA3	MG, 48 MG, 6 MG19
ACTEMRA ACTPEN3	AUSTEDO XR PATIENT TITRATION
ACTIMMUNE4	ORAL TABLET EXTENDED
acyclovir sodium intravenous solution 50	RELEASE THERAPY PACK 12 & 18 &
mg/ml	24 & 30 MG, 6 & 12 & 24 MG19
adalimumab-adbm (2 pen) 5, 6	AUVELITY20
adalimumab-adbm (2 syringe) subcutaneous	AYVAKIT21
prefilled syringe kit 10 mg/0.2ml, 20	azathioprine oral tablet 50 mg 285
mg/0.4ml, 40 mg/0.8ml	azithromycin intravenous solution
adalimumab-adbm(cd/uc/hs strt) 5, 6	reconstituted 500 mg285
adalimumab-adbm(ps/uv starter) 5, 6	В
adapalene238	BALVERSA ORAL TABLET 3 MG, 4
adefovir dipivoxil88	MG, 5 MG22
ADEMPAS7	BARACLUDE 88
AKEEGA9	BENLYSTA24
albuterol sulfate inhalation nebulization	benztropine mesylate96
solution (2.5 mg/3ml) 0.083%, 0.63	BESREMI25
mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml . 285	bexarotene
ALECENSA10	bosentan
alosetron hcl11	BOSULIF ORAL CAPSULE 100 MG, 50
alprazolam oral tablet 0.25 mg, 0.5 mg, 1	MG29
mg, 2 mg97, 98	BOSULIF ORAL TABLET 100 MG, 400
ALUNBRIG12	MG, 500 MG29
ambrisentan13	BRAFTOVI30
	BRIVIACT31

Prior Authorization Criteria

BRONCHITOL 32	colistimethate sodium (cba) injection
BRUKINSA33	solution reconstituted 150 mg 285
budesonide inhalation suspension 0.25	COMETRIQ (100 MG DAILY DOSE) 43
mg/2ml, 0.5 mg/2ml, 1 mg/2ml 285	COMETRIQ (140 MG DAILY DOSE) 43
butalbital-acetaminophen90	COMETRIQ (60 MG DAILY DOSE) 43
butalbital-apap-caffeine 90	COPIKTRA 44
BYDUREON BCISE86	COSENTYX (300 MG DOSE)45
C	COSENTYX SENSOREADY (300 MG) 45
CABOMETYX 34	COSENTYX UNOREADY45
calcitonin (salmon) nasal solution 200	COTELLIC46
unit/act285	cromolyn sodium inhalation nebulization
calcitriol oral capsule 0.25 mcg, 0.5 mcg285	solution 20 mg/2ml285
calcitriol oral solution 1 mcg/ml 285	cyclobenzaprine hcl104
CALQUENCE35	CYCLOPHOSPHAMIDE ORAL
CAMZYOS36	CAPSULE 25 MG 285
CAPLYTA17	cyclophosphamide oral capsule 50 mg 285
CAPRELSA ORAL TABLET 100 MG, 300	cyclophosphamide oral tablet 25 mg 285
MG37	CYCLOPHOSPHAMIDE ORAL TABLET
carglumic acid38	50 MG285
CAYSTON 39	cyclosporine modified oral capsule 100 mg,
cefoxitin sodium intravenous solution	25 mg, 50 mg
reconstituted 1 gm, 10 gm, 2 gm 285	cyclosporine modified oral solution 100
cefuroxime sodium injection solution	mg/ml285
reconstituted 750 mg285	cyclosporine oral capsule 100 mg, 25 mg
cefuroxime sodium intravenous solution	285
reconstituted 1.5 gm 285	CYSTAGON47
chlordiazepoxide hcl97, 98	D
chlordiazepoxide-amitriptyline94	dalfampridine er48
chlordiazepoxide-clidinium92	DAURISMO 49
CIMZIA40	DAYBUE 50
CIMZIA (2 SYRINGE)40	deferasirox51
cinacalcet hcl oral tablet 30 mg, 60 mg, 90	deferiprone52
mg285	DEXTROSE-SODIUM CHLORIDE
ciprofloxacin in d5w intravenous solution	INTRAVENOUS SOLUTION 10-0.2 %,
200 mg/100ml285	10-0.45 %, 2.5-0.45 %
clindamycin phosphate injection solution	DIACOMIT53
900 mg/6ml 285	diazepam intensol 101, 102
clobazam oral suspension41	diazepam oral solution 101, 102
clobazam oral tablet41	diazepam oral tablet101, 102
clonazepam oral tablet 0.5 mg, 1 mg, 2 mg	diclofenac epolamine 54
101, 102	DIGOXIN ORAL SOLUTION91
clonazepam oral tablet dispersible 0.125 mg,	digoxin oral tablet 250 mcg91
0.25 mg, 0.5 mg, 1 mg, 2 mg 101, 102	dimethyl fumarate55
clorazepate dipotassium 101, 102	dimethyl fumarate starter pack 55

doxy 100 intravenous solution reconstituted	FIRMAGON76
100 mg285	FIRMAGON (240 MG DOSE)76
dronabinol56	fluconazole in sodium chloride intravenous
droxidopa57	solution 200-0.9 mg/100ml-%, 400-0.9
DUPIXENT 58, 59	mg/200ml-%285
E	FOTIVDA77
ELIGARD SUBCUTANEOUS KIT 22.5	FRUZAQLA78
MG, 30 MG, 45 MG, 7.5 MG 139	FULPHILA79
EMGALITY60	FYCOMPA80
EMGALITY (300 MG DOSE)60	G
EMSAM 61	GAMMAGARD INJECTION SOLUTION
ENBREL 62, 63	2.5 GM/25ML285
ENBREL MINI 62, 63	GAMMAGARD S/D LESS IGA
ENBREL SURECLICK 62, 63	INTRAVENOUS SOLUTION
ENGERIX-B INJECTION SUSPENSION	RECONSTITUTED 10 GM, 5 GM 286
20 MCG/ML285	GAMMAPLEX INTRAVENOUS
ENGERIX-B INJECTION SUSPENSION	SOLUTION 10 GM/100ML, 10
PREFILLED SYRINGE 10 MCG/0.5ML,	GM/200ML, 20 GM/200ML, 5
20 MCG/ML285	GM/50ML286
ENSPRYNG 64	GAMUNEX-C INJECTION SOLUTION 1
entecavir 88	GM/10ML286
ENVARSUS XR ORAL TABLET	GAVRETO81
EXTENDED RELEASE 24 HOUR 0.75	gefitinib82
MG, 1 MG, 4 MG285	GILOTRIF83
EPIDIOLEX65	glatiramer acetate84
ergoloid mesylates93	GLEOSTINE85
ERIVEDGE 67	granisetron hcl oral tablet 1 mg 286
ERLEADA 68	guanfacine hcl er99
erlotinib hcl69	H
estazolam97, 98	heparin sodium (porcine) injection solution
everolimus oral tablet 0.25 mg, 0.5 mg, 0.75	10000 unit/ml, 20000 unit/ml, 5000
mg, 1 mg285	unit/ml286
everolimus oral tablet 10 mg, 2.5 mg, 5 mg,	HEPLISAV-B INTRAMUSCULAR
7.5 mg8	<b>SOLUTION PREFILLED SYRINGE 20</b>
everolimus oral tablet soluble 2 mg, 3 mg, 5	MCG/0.5ML286
mg70	HUMIRA (2 PEN) SUBCUTANEOUS
EVRYSDI71	PEN-INJECTOR KIT 40 MG/0.4ML, 40
F	MG/0.8ML, 80 MG/0.8ML 105, 106
FANAPT 17	HUMIRA (2 SYRINGE)
FANAPT TITRATION PACK 17	SUBCUTANEOUS PREFILLED
FASENRA72	SYRINGE KIT 10 MG/0.1ML, 20
FASENRA PEN72	MG/0.2ML, 40 MG/0.4ML, 40
febuxostat	MG/0.8ML105, 106
fentanyl citrate74	HUMIRA-CD/UC/HS STARTER 105, 106
FILSPARI75	

HUMIRA-PED>/=40KG UC STARTER	ISOLYTE-P IN D5W INTRAVENOUS
105, 106	SOLUTION286
HUMIRA-PSORIASIS/UVEIT STARTER	ISOLYTE-S PH 7.4 INTRAVENOUS
105, 106	SOLUTION286
hydroxyzine hcl95	itraconazole123
hydroxyzine pamoate95	ivabradine hcl124
HYFTOR107	ivermectin125
I	IWILFIN126
IBRANCE 108	J
icatibant acetate109	JAKAFI127
ICLUSIG ORAL TABLET 10 MG, 15 MG,	JAYPIRCA ORAL TABLET 100 MG, 50
30 MG, 45 MG110	MG128
IDHIFA ORAL TABLET 100 MG, 50 MG	JYLAMVO ORAL SOLUTION 2 MG/ML
111	286
imatinib mesylate oral tablet 100 mg, 400	K
mg112	KALYDECO129
IMBRUVICA ORAL CAPSULE 140 MG,	kcl in dextrose-nacl intravenous solution 10-
70 MG 113	5-0.45 meg/l-%-%, 20-5-0.2 meg/l-%-%,
IMBRUVICA ORAL SUSPENSION 113	20-5-0.45 meg/l-%-%, 20-5-0.9 meg/l-%-
IMBRUVICA ORAL TABLET 113	%, 30-5-0.45 meq/l-%-%, 40-5-0.45
IMOVAX RABIES INTRAMUSCULAR	meq/l-%-%286
SUSPENSION RECONSTITUTED 2.5	KCL IN DEXTROSE-NACL
UNIT/ML286	<b>INTRAVENOUS SOLUTION 40-5-0.9</b>
IMPAVIDO114	MEQ/L-%-%286
INCRELEX115	KCL-LACTATED RINGERS-D5W
INLYTA117	INTRAVENOUS SOLUTION 20 MEQ/L
INQOVI118	286
INREBIC119	KERENDIA130
INVEGA HAFYERA INTRAMUSCULAR	KISQALI (200 MG DOSE) 131
SUSPENSION PREFILLED SYRINGE	KISQALI (400 MG DOSE) 131
1092 MG/3.5ML, 1560 MG/5ML 120	KISQALI (600 MG DOSE)131
INVEGA SUSTENNA	KISQALI FEMARA (200 MG DOSE) 132
INTRAMUSCULAR SUSPENSION	KISQALI FEMARA (400 MG DOSE) 132
PREFILLED SYRINGE 117	KISQALI FEMARA (600 MG DOSE) 132
MG/0.75ML, 156 MG/ML, 234	KORLYM133
MG/1.5ML, 39 MG/0.25ML, 78	KOSELUGO134
MG/0.5ML 121	KRAZATI 135
INVEGA TRINZA INTRAMUSCULAR	L
SUSPENSION PREFILLED SYRINGE	lapatinib ditosylate136
273 MG/0.88ML, 410 MG/1.32ML, 546	ledipasvir-sofosbuvir89
MG/1.75ML, 819 MG/2.63ML 122	lenalidomide137
ipratropium bromide inhalation solution	LENVIMA (10 MG DAILY DOSE) 138
0.02 %	LENVIMA (12 MG DAILY DOSE) 138
ipratropium-albuterol inhalation solution	LENVIMA (14 MG DAILY DOSE) 138
0.5-2.5 (3) mg/3ml	LENVIMA (18 MG DAILY DOSE) 138

LENVIMA (20 MG DAILY DOSE) 138	miglustat155	
LENVIMA (24 MG DAILY DOSE) 138	modafinil42	
LENVIMA (4 MG DAILY DOSE) 138	MOUNJARO86	
LENVIMA (8 MG DAILY DOSE) 138	multiple electro type 1 ph 5.5 intravenous	
leuprolide acetate139	solution286	
LEUPROLIDE ACETATE (3 MONTH)139	mycophenolate mofetil oral capsule 250 mg	
1-glutamine		
lidocaine external ointment	mycophenolate mofetil oral suspension	
lidocaine hcl141	reconstituted 200 mg/ml286	
linezolid intravenous	mycophenolate mofetil oral tablet 500 mg	
linezolid oral suspension reconstituted 142		
linezolid oral tablet142	mycophenolate sodium oral tablet delayed	
LIVTENCITY 143	release 180 mg, 360 mg 286	
LONSURF144	N	
lorazepam 101, 102	NAYZILAM 156	
lorazepam intensol 101, 102	NERLYNX 157	
LORBRENA ORAL TABLET 100 MG, 25	NINLARO158	
MG145	NUBEQA159	
LUMAKRAS ORAL TABLET 120 MG,	NUEDEXTA160	
320 MG146	NUPLAZID161	
LUPRON DEPOT (1-MONTH) 139	0	
LUPRON DEPOT (3-MONTH) 139	octreotide acetate162	
LUPRON DEPOT (4-MONTH) 139	ODOMZO163	
LUPRON DEPOT (6-MONTH) 139	OFEV164	
LYBALVI17	OGSIVEO ORAL TABLET 100 MG, 150	
LYNPARZA 147, 148	MG, 50 MG165	
LYTGOBI (12 MG DAILY DOSE) 149	OJEMDA ORAL SUSPENSION	
LYTGOBI (16 MG DAILY DOSE) 149	RECONSTITUTED166	
LYTGOBI (20 MG DAILY DOSE) 149	OJEMDA ORAL TABLET 166	
M	OJJAARA167	
MATULANE 150	OMNITROPE87	
MAVYRET 89	ondansetron hcl oral solution 4 mg/5ml. 286	
MAYZENT ORAL TABLET 0.25 MG, 1	ondansetron hcl oral tablet 4 mg, 8 mg 286	
MG, 2 MG151	ondansetron oral tablet dispersible 4 mg, 8	
MAYZENT STARTER PACK 151	mg286	
megestrol acetate	ONUREG168	
MEKINIST 152	ORENCIA169	
MEKTOVI153	ORENCIA CLICKJECT169	
methotrexate sodium (pf) injection solution	ORGOVYX170	
50 mg/2ml	ORKAMBI 171	
methotrexate sodium injection solution 50	orphenadrine citrate er104	
mg/2ml 286	ORSERDU ORAL TABLET 345 MG, 86	
metronidazole intravenous solution 500	MG172	
mg/100ml286	OTEZLA ORAL TABLET173	
mifepristone 154		

OTEZLA ORAL TABLET THERAPY	PREHEVBRIO INTRAMUSCULAR		
PACK 173	SUSPENSION 10 MCG/ML 286		
OZEMPIC (0.25 OR 0.5 MG/DOSE) 86	PREVYMIS 183		
OZEMPIC (1 MG/DOSE)	PROGRAF ORAL PACKET 0.2 MG, 1 MG		
SUBCUTANEOUS SOLUTION PEN-	286		
INJECTOR 2 MG/1.5ML, 4 MG/3ML 86	PROLASTIN-C INTRAVENOUS		
OZEMPIC (2 MG/DOSE)86	SOLUTION 1000 MG/20ML 286		
P	PROLIA 184		
PANRETIN 174	PROMACTA ORAL PACKET 12.5 MG,		
paricalcitol oral capsule 1 mcg, 2 mcg, 4	25 MG 185		
mcg286	PROMACTA ORAL TABLET 185		
pazopanib hcl	promethazine hcl95		
PEGASYS 176	promethegan95		
PEMAZYRE177	PULMOZYME 186		
penicillamine178	PURIXAN187		
penicillin g potassium injection solution	pyrimethamine188		
reconstituted 20000000 unit 286	Q		
penicillin g sodium injection solution	QINLOCK 189		
reconstituted 5000000 unit 286	quinine sulfate		
pentamidine isethionate inhalation solution	Ŕ		
reconstituted 300 mg 286	RABAVERT INTRAMUSCULAR		
pentamidine isethionate injection solution	SUSPENSION RECONSTITUTED 287		
reconstituted 300 mg	RAVICTI191		
PERSERIS SUBCUTANEOUS	RECOMBIVAX HB INJECTION		
PREFILLED SYRINGE 120 MG, 90 MG	SUSPENSION 10 MCG/ML, 40		
286	MCG/ML, 5 MCG/0.5ML287		
PIQRAY (200 MG DAILY DOSE) 179	RECOMBIVAX HB INJECTION		
PIQRAY (250 MG DAILY DOSE) 179	SUSPENSION PREFILLED SYRINGE		
PIQRAY (300 MG DAILY DOSE) 179	10 MCG/ML, 5 MCG/0.5ML 287		
pirfenidone180	REGRANEX192		
POMALYST181	REPATHA193		
posaconazole182	REPATHA PUSHTRONEX SYSTEM. 193		
POTASSIUM CHLORIDE IN NACL	REPATHA SURECLICK 193		
<b>INTRAVENOUS SOLUTION 20-0.45</b>	RETACRIT66		
MEQ/L-%, 40-0.9 MEQ/L-% 286	RETEVMO ORAL CAPSULE 40 MG, 80		
potassium chloride in nacl intravenous	MG194		
solution 20-0.9 meq/l-%286	RETEVMO ORAL TABLET 120 MG, 160		
POTASSIUM CHLORIDE	MG, 40 MG, 80 MG194		
INTRAVENOUS SOLUTION 10	REVLIMID195		
MEQ/100ML, 20 MEQ/100ML, 40	REXULTI17		
MEQ/100ML286	REZLIDHIA196		
potassium chloride intravenous solution 2	REZUROCK197		
meq/ml, 2 meq/ml (20 ml)286	riluzole198		
potassium cl in dextrose 5% intravenous	RINVOQ199		
solution 20 meq/l286	RINVOQ LQ200		

ROZLYTREK ORAL CAPSULE 100 MG,	T	
200 MG201	TABRECTA222	
ROZLYTREK ORAL PACKET 201	tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg	
RUBRACA202	287	
rufinamide oral suspension	tadalafil223	
rufinamide oral tablet	TAFINLAR224	
RYBELSUS86	TAGRISSO225	
RYDAPT204	TAKHZYRO SUBCUTANEOUS	
S	SOLUTION226	
sapropterin dihydrochloride 205	TAKHZYRO SUBCUTANEOUS	
SCEMBLIX206	SOLUTION PREFILLED SYRINGE 150	
SECUADO 17	MG/ML, 300 MG/2ML 226	
SIGNIFOR207	TALZENNA227	
sildenafil citrate oral tablet 20 mg 208	TASIGNA228	
sirolimus oral solution 1 mg/ml 287	TAVNEOS229	
sirolimus oral tablet 0.5 mg, 1 mg, 2 mg 287	tazarotene external cream	
SIRTURO	tazarotene external gel230	
SKYRIZI (150 MG DOSE)210	TAZORAC	
SKYRIZI PEN210	TAZVERIK231	
SKYRIZI SUBCUTANEOUS SOLUTION	TEFLARO INTRAVENOUS SOLUTION	
CARTRIDGE 180 MG/1.2ML, 360	RECONSTITUTED 400 MG, 600 MG	
MG/2.4ML210	287	
SKYRIZI SUBCUTANEOUS SOLUTION	temazepam97, 98	
PREFILLED SYRINGE210	TEPMETKO	
sodium oxybate	TERIPARATIDE (RECOMBINANT) 233	
sofosbuvir-velpatasvir	testosterone cypionate	
SOLTAMOX212	testosterone enanthate	
SOMAVERT SUBCUTANEOUS	testosterone transdermal gel 10 mg/act (2%),	
SOLUTION RECONSTITUTED 10 MG,	12.5 mg/act (1%), 20.25 mg/1.25gm	
15 MG, 20 MG, 25 MG, 30 MG 213	(1.62%), 20.25 mg/act (1.62%), 25	
sorafenib tosylate214	mg/2.5gm (1%), 40.5 mg/2.5gm (1.62%),	
SPRYCEL ORAL TABLET 100 MG, 140	50 mg/5gm (1%), 40.5 mg/2.5gm (1.02%),	
MG, 20 MG, 50 MG, 70 MG, 80 MG 215	testosterone transdermal solution	
STELARA SUBCUTANEOUS	tetrabenazine oral tablet 12.5 mg, 25 mg 234	
SOLUTION216, 217	THALOMID	
STELARA SUBCUTANEOUS	TIBSOVO	
SOLUTION PREFILLED SYRINGE 45	TIGECYCLINE INTRAVENOUS	
MG/0.5ML, 90 MG/ML 216, 217	SOLUTION RECONSTITUTED 50 MG	
STIVARGA218	287	
sunitinib malate	TOBI PODHALER237	
SYMLINPEN 120		
SYMLINPEN 60	tobramycin inhalation nebulization solution	
SYMPAZAN	300 mg/5ml	
SYNAREL221	tobramycin sulfate injection solution 10	
STNAKEL221	mg/ml, 80 mg/2ml	
	TOTE OTHER COLLARS 740	

TPN ELECTROLYTES INTRAVENOUS	VUMERITY 264	
CONCENTRATE 287	W	
TRELSTAR MIXJECT	WELIREG265	
INTRAMUSCULAR SUSPENSION	X	
RECONSTITUTED 11.25 MG, 22.5 MG,	XALKORI ORAL CAPSULE266	
3.75 MG241	XALKORI ORAL CAPSULE SPRINKLE	
TREMFYA 242	150 MG, 20 MG, 50 MG266	
tretinoin external238	XCOPRI (250 MG DAILY DOSE) 267	
trientine hcl	XCOPRI (350 MG DAILY DOSE) 267	
trihexyphenidyl hcl96	XCOPRI ORAL TABLET 100 MG, 150	
TRIKAFTA244	MG, 200 MG, 25 MG, 50 MG 267	
TRULICITY 86	XCOPRI ORAL TABLET THERAPY	
TRUQAP245	PACK267	
TUKYSA246	XDEMVY268	
TURALIO247	XELJANZ ORAL SOLUTION269	
TYMLOS248	XELJANZ ORAL TABLET 10 MG, 5 MG	
U	269	
UBRELVY249	XELJANZ XR269	
V	XERMELO270	
VALCHLOR250	XGEVA271	
VALTOCO 20 MG DOSE251	XOLAIR272	
VANFLYTA252	XOSPATA273	
VEMLIDY88	XPOVIO (100 MG ONCE WEEKLY) 274	
VENCLEXTA253	XPOVIO (40 MG ONCE WEEKLY) 274	
VENCLEXTA STARTING PACK 253	XPOVIO (40 MG TWICE WEEKLY) 274	
VEOZAH254	XPOVIO (60 MG ONCE WEEKLY) 274	
VERQUVO255	XPOVIO (60 MG TWICE WEEKLY) 274	
VERSACLOZ 17	XPOVIO (80 MG ONCE WEEKLY) 274	
VERZENIO256	XPOVIO (80 MG TWICE WEEKLY) 274	
vigabatrin257	XTANDI275	
vigadrone257	Υ	
vigpoder	YONSA276	
VIJOICE258	Z	
VITRAKVI ORAL CAPSULE 100 MG, 25	zaleplon 103	
MG259	ZARXIO277	
VITRAKVI ORAL SOLUTION 259	ZEJULA278	
VIZIMPRO260	ZELBORAF279	
VONJO261	ziprasidone mesylate intramuscular solution	
VORICONAZOLE INTRAVENOUS 262	reconstituted 20 mg287	
voriconazole oral suspension reconstituted	ZOLINZA280	
262	zolpidem tartrate103	
voriconazole oral tablet 200 mg, 50 mg. 262	ZTALMY281	
VOWST263	ZURZUVAE ORAL CAPSULE 20 MG, 25	
VRAYLAR ORAL CAPSULE 1.5 MG, 3	MG, 30 MG282	
MG, 4.5 MG, 6 MG 17		

ZYDELIG ORAL T	CABLET 100 MG, 150	ZYKADIA	284
MG	283		